







## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07D 405/14, 405/12, 401/14, 209/20, A61K 31/495, 31/445</b>		<b>A1</b>	(11) International Publication Number: <b>WO 99/67237</b>
			(43) International Publication Date: 29 December 1999 (29.12.99)
(21) International Application Number: PCT/DK99/00326 (22) International Filing Date: 14 June 1999 (14.06.99) (30) Priority Data: PA 1998 00820      19 June 1998 (19.06.98)      DK 60/092,823      14 July 1998 (14.07.98)      US (71) Applicant (for all designated States except US): H. LUNDBECK A/S [DK/DK]; Ottiliavej 9, DK-2500 Valby-Copenhagen (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): MOLTZEN, Ejner, Knud [DK/DK]; Toftekærsvvej 29, DK-2820 Gentofte (DK). MIKKELSEN, Ivan [DK/DK]; Strandvejen 97, DK-4600 Køge (DK). KROG-JENSEN, Christian [SE/DK]; Hornemansgade 23, DK-2100 København Ø (DK). (74) Common Representative: H. LUNDBECK A/S; Petersen, John, Meidahl, Ottiliavej 9, DK-2500 Valby-Copenhagen (DK).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: 4,5,6 AND 7-INDOLE AND INDOLINE DERIVATIVES, THEIR PREPARATION AND USE



## (57) Abstract

A substituted 4-, 5-, 6-, and 7-indole or indoline derivative of Formula (I) wherein A is a group having formula (IIA), (IIB), (IIC) wherein X, U, Y, R<sup>3</sup> to R<sup>12</sup>, Z, W, n and m are as defined above. The compounds are potent serotonin reuptake inhibitors and have 5-HT<sub>1A</sub> receptor antagonistic activity.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

#### 4, 5, 6 and 7-indole and indoline derivatives, their preparation and use

The present invention relates to novel 4, 5, 6 and 7-indole and indoline derivatives  
5 which are potent serotonin reuptake inhibitors, pharmaceutical compositions  
containing these compounds and the use thereof for the treatment of disorders or  
diseases responsive to the inhibition of serotonin re-uptake. The compounds of the  
invention also possess antagonistic activity at 5-HT<sub>1A</sub> receptors and are considered to  
be particularly useful for the treatment of depression.

10

#### Background

Selective serotonin (or 5-HT) reuptake inhibitors (SSRI's) such as fluoxetine,  
paroxetine, sertraline, fluvoxamine and citalopram represent a major step forward in  
15 the treatment of depression because they have fewer and less severe side effects  
compared to first generation antidepressant (tricyclics and non-selective MAO  
inhibitors). The side effects associated with first generation antidepressants are such  
that they cause some patients to withdraw from treatment.

20 SSRI's and all other antidepressants currently available suffer from a serious  
drawback in that several weeks of treatment is necessary to produce the therapeutic  
effect. The late onset of action is a significant problem, particularly in the treatment  
of patients with severe depression and suicide potential. Further, one in three patients  
are not responsive to SSRI's.

25

Electrophysiological experiments in rats have shown that acute administration of  
SSRI's reduces firing of 5-HT neurons of dorsal raphe nucleus in the rodent brain,  
whereas sustained treatment with SSRI's leads to normalization of the firing activity  
of the 5-HT neurons (Arborelius, L. *et al*, *Naunyn-Schmiedeberg's Arch. Pharmacol.*  
30 *1995*, 352, 157; Gartside, S.E. *et al*, *Br. J. Pharmacol.* *1995*, 115, 1064; Chaput, Y. *et*  
*al*, *Naunyn-Schmiedeberg's Arch. Pharmacol.* *1986*, 33, 342).

CONFIRMATION COPY

Further, it has been shown that the recovery of the firing activity of 5-HT neurons is linked to desensitization of somatodendritic 5-HT<sub>1A</sub> autoreceptors (Le Poul, E. *et al*, *Naunyn-Schmiedeberg's Arch. Pharmacol.* **1995**, 352, 141; Invernizzi, R. *et al*, *Eur. J. Pharmacol.* **1994**, 260, 243).

5

It has thus been suggested that simultaneous administration of SSRI's and an agent causing rapid desensitization or inhibition of the 5-HT<sub>1A</sub> receptor mediated feed back mechanism would lead to rapid onset of antidepressive effect (Artigas, F. *et al*, *Trends Neurosci.* **1996**, 19, 378; De Vry, J., *et al*, *Drug News Perspec.* **1996**, 9, 270).

10

The effect of combined administration of a compound that inhibits serotonin reuptake and a 5-HT<sub>1A</sub> receptor antagonist has been evaluated in several studies (Innis, R.B. *et al*, *Eur. J. Pharmacol.*, **1987**, 143, p 195-204 and Gartside, S.E., *Br. J. Pharmacol.* **1995**, 115, p 1064-1070, Blier, P. *et al*, *Trends Pharmacol. Sci.* **1994**, 15, 220). In these studies it was found that 5-HT<sub>1A</sub> receptor antagonists inhibit the decrease in firing caused by acute administration of serotonin reuptake inhibitors.

15

Further, treatment with a combination of pindolol (a well known 5-HT<sub>1A</sub> receptor and  $\beta$ -adrenoceptor antagonist) and SSRI's has been evaluated in clinical trials.

20


A remarkable improvement of the mood of patients was reported within one week. In addition, combined administration of pindolol and a SSRI was shown to have a good effect on patients who were non-responsive to treatment with currently available antidepressants (Artigas F. *et al*, *Arch. Gen. Psychiatry*, **1994**, 51, p 248-251 and Blier, P. *et al*, *J. Clin. Psychopharmacol.* **1995**, 15, p 217-222).

25

Several patent applications have been filed which cover the use of a combination of a 5-HT<sub>1A</sub> antagonist and a serotonin reuptake inhibitor for the treatment of depression (see EP-A2-687 472 and EP-A2-714 663).


30

DE patent application No. 4414113 discloses certain 4-(indol-3-yl)-1-(indol-3-yl-alkylene)-piperidines having the general formula



wherein n is 2-6 and the other substituents are as defined in the application. The compounds herein are claimed to have serotonin antagonistic and agonistic activities and to have effect on DOPA-accumulation in striatum and 5-HTP accumulation in N. Raphe. No biological data are given.


WO patent publication No. 94/21626 discloses compounds having the general formula



wherein  $R^2$  is heteroaryl and the other substituents are as defined in the application. A compound wherein  $R^2$  is 5-indolyl which is structurally closely related to the compounds of the invention is specifically mentioned herein. No data are given. The compounds are only said to give  $K_i$  values of less than 1.5  $\mu\text{M}$  in a test for displacement of  $^3\text{H}$  spiperone from human dopamine  $D_4$  receptor subtypes in clonal cell lines. WO patent publication No. 94/21627 and No. 94/21630 relate to similar compounds having affinity for human dopamine  $D_4$  receptors.

WO patent publication No. 95/33721 relates to 1-(indanemethyl, dihydrobenzofuranylmethyl, dihydrobenzothiophenylmethyl)piperidine, tetrahydro-pyridine, or

piperazine derivatives having the general formula



- wherein one of X and Y is CH<sub>2</sub> and the other is selected from the group consisting of CH<sub>2</sub>, O, or S, Ar is aryl or heteroaryl, e.g. 1-, 2-, or 3-indolyl and the other substituents is as defined in the application. The compounds interact with central
- 5 5-HT receptors, in particular with 5-HT<sub>1A</sub> and 5-HT<sub>2A</sub> receptors. Some of the compounds are said to have 5-HT reuptake inhibiting effect.


### Object of the Invention

- 10 It is the object of the present invention to provide compounds with potent serotonin reuptake inhibiting activity as well as antagonistic properties at 5-HT<sub>1A</sub> receptors. Such compounds may be useful as fast onset of action medicaments for the treatment of affective disorders, such as depression.
- 15 A further object of the present invention is to provide a pharmaceutical composition comprising these compounds as active ingredients.

### Summary of the Invention


- 20 The invention then, inter alia, comprises the following alone or in combination:

A substituted 4-, 5-, 6-, or 7-indole or indoline derivative of formula (I)



wherein W is N, C, CH or COH and the dotted lines indicate optional bonds and

5 wherein A is a group having the formula



10 wherein X is CR<sup>1A</sup>, CHR<sup>1A</sup>, N, NR<sup>1B</sup>, O, or S, where R<sup>1A</sup> is as defined for R<sup>3</sup> to R<sup>9</sup> below, and where R<sup>1B</sup> is as defined for R<sup>10</sup> below;


Y is CR<sup>2A</sup>, CHR<sup>2A</sup>, N, NR<sup>2B</sup>, O, or S, where R<sup>2A</sup> is as defined for R<sup>3</sup> to R<sup>9</sup> below and where R<sup>2B</sup> is as defined for R<sup>10</sup> below, and

15 the dotted lines indicate optional bonds;

provided that X and Y are not both O or S;

A is a group having the formula

20




wherein X is CR<sup>1A</sup>, CHR<sup>1A</sup>, N, NR<sup>1B</sup>, O, or S, where R<sup>1A</sup> is as defined for R<sup>3</sup> to R<sup>9</sup> below, and where R<sup>1B</sup> is as defined for R<sup>10</sup> below;

5

U is C, CH, or N; and

the dotted lines indicate optional bonds;

10 A is a group having the formula



wherein U is C, CH, or N;

15

Y is CR<sup>2A</sup>, CHR<sup>2A</sup>, N, NR<sup>2B</sup>, O, or S, where R<sup>2A</sup> is as defined for R<sup>3</sup> to R<sup>9</sup> below and where R<sup>2B</sup> is as defined for R<sup>10</sup> below;

and the dotted lines indicate optional bonds;

20

n is 0, 1, 2, 3, 4, or 5, and m is 0, 1, 2, 3, 4, or 5;


Z is CH<sub>2</sub>, O, S, CO, SO, or SO<sub>2</sub>, provided that if n is 0 then Z is CH<sub>2</sub>;



- $R^3$ - $R^9$  and  $R^{11}$  to  $R^{12}$  are independently selected from hydrogen, halogen, cyano, nitro,  $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$  alkoxy,  $C_{1-6}$ -alkylthio, hydroxy, hydroxy- $C_{1-6}$ -alkyl,  $C_{1-6}$ -alkoxycarbonyl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl,  $C_{1-6}$ -alkylcarbonyl, phenylcarbonyl, halogen substituted phenylcarbonyl, trifluoromethyl, trifluoromethylsulfonyloxy,  $C_{1-6}$  alkylsulfonyl, aryl and heteroaryl, and/or two adjacent groups taken from  $R^3$  -  $R^9$  may together form a methylenedioxy group, and/or two adjacent groups  $R^7$  -  $R^9$  may together form a cyclopentyl or cyclohexyl ring which may be substituted with one or more methyl groups,
- 10 and/or one of  $R^3$ - $R^9$  may alternatively be a group  $-NR^{13}R^{14}$  wherein  $R^{13}$  is as defined for  $R^{10}$  below and  $R^{14}$  is hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$  alk(en/yn)yl, aryl, heteroaryl, aryl- $C_{1-6}$  alkyl, or heteroaryl- $C_{1-6}$ -alkyl;
- 15  $R^{10}$  is
- hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, aryl, heteroaryl, aryl- $C_{1-6}$ -alkyl, heteroaryl- $C_{1-6}$ -alkyl, acyl, thioacyl,  $C_{1-6}$ -alkylsulfonyl, trifluoromethylsulfonyl, arylsulfonyl, or heteroarylsulfonyl;
  - $R^{15}VCO$ - wherein V is O or S and  $R^{15}$  is  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, aryl, or heteroaryl; or
  - a group  $R^{16}R^{17}NCO$ - or  $R^{16}R^{17}NCS$ - wherein  $R^{16}$  and  $R^{17}$  are independently selected from hydrogen,  $C_{1-6}$ -alk(en/yn)yl,  $C_{3-8}$ -cycloalk(en)yl,  $C_{3-8}$ -cycloalk(en)yl- $C_{1-6}$ -alk(en/yn)yl, heteroaryl, or aryl, or  $R^{16}$  and  $R^{17}$  together with the N-atom to which they are linked, form a pyrrolidinyl, piperidinyl, morpholinyl, or
- 25 perhydroazepin group;

or an acid addition salt thereof.

- In a particular embodiment, the compounds of the invention are compounds wherein
- 30 A is a group of formula (IIA) including such compounds wherein A is a group having the formulas illustrated below:




or

wherein  $R^3$  to  $R^6$  and the dotted lines are as defined above.


5

In a particular embodiment A is a group having the formula



wherein  $R^3$  to  $R^6$  and the dotted lines are as defined above.

- 5 In another particular embodiment the present invention relates to compounds having the formula




- 10 wherein  $R^7$  to  $R^{12}$ , W, A, Z, n, m and the dotted lines are as defined above.

In a particular embodiment Z is  $\text{CH}_2$  and  $n + m$  is 0, 1, 2, 3, 4, 5, or 6.


- In another embodiment the present invention relates to compounds having the  
15 formula II above and A is a group having the formula IIA above.

In another particular embodiment the present invention relates to compounds having the formula II above and A is a group having the formula



wherein  $R^3$  to  $R^6$  and the dotted lines are as defined above.

- 10 In a further particular embodiment the present invention relates to compounds having the formula II above and A is a group having the formula



wherein  $R^3$  to  $R^6$  and the dotted lines are as defined above.

5

In a particular embodiment the compounds of the invention are the compounds wherein  $R^3$ - $R^9$  and  $R^{11}$ - $R^{12}$  is hydrogen, halogen, cyano, nitro,  $C_{1-6}$ -alkyl,  $C_{1-6}$ -alkoxy hydroxy, hydroxy- $C_{1-6}$ -alkyl,  $C_{1-6}$ -alkoxycarbonyl and trifluoromethyl; and  $R^{10}$  is hydrogen.

10

In another particular embodiment of the invention, W is N.

Examples of compounds according to the invention are the compounds

1-(2-(3-Benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,

15 1-(3-Benzofuranylmethyl)-4-(1H-indol-4-yl)piperazine,

1-(2-(5-Fluoro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,

1-(4-(5-Fluoro-3-benzofuranyl)-1-butyl)-4-(1H-indol-4-yl)piperazine,

1-(2-(1H-Indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,

1-(3-(1H-Indol-3-yl)-1-propyl)-4-(1H-indol-4-yl)piperazine,

20 1-(4-(1H-Indol-3-yl)-1-butyl)-4-(1H-indol-4-yl)piperazine,

1-(3-(5-Fluoro-3-benzofuranyl)-1-propyl)-4-(1H-indol-4-yl)piperazine,

- 1-(2-(2-Methyl-4,5,6,7-tetrafluoro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(3-Indazolyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-3-indazolyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(7-Cyano-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
5 1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(4-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)-1,2,3,6-tetrahydropyridine,  
1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)-1,2,3,6-tetrahydropyridine,  
10 1-(2-(7-Bromo-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(1-Allyl-1H-indol-4-yl)-4-(2-(6-chloro-1H-indol-3-yl)ethyl)piperazine,  
1-(1-Allyl-1H-indol-4-yl)-4-(2-(5-fluoro-1H-indol-3-yl)ethyl)piperazine,  
1-(1-Benzyl-1H-indol-4-yl)-4-(2-(6-chloro-1H-indol-3-yl)ethyl)piperazine,  
1-(1-Benzyl-1H-indol-4-yl)-4-(2-(5-fluoro-1H-indol-3-yl)ethyl)piperazine,  
15 1-(1-Benzyl-1H-indol-4-yl)-4-(2-(5-bromo-1H-indol-3-yl)ethyl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine,  
1-(2-(1H-Indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine,  
1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine,  
1-(2-(5-Bromo-1H-indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine,  
20 1-(1-Benzyl-1H-indol-4-yl)-4-(2-(1H-indol-3-yl)ethyl)piperazine,  
1-(2-(5-Bromo-1H-indol-3-yl)ethyl)-4-(1H-indol-5-yl)piperazine,  
1-(2-(5-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-5-yl)piperazine,  
1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(6-hydroxymethyl-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-hydroxymethyl-1H-indol-4-yl)piperazine,  
25 1-(2-(5-Bromo-1H-indol-3-yl)ethyl)-4-(6-hydroxymethyl-1H-indol-4-yl)piperazine,  
1-(3-(6-Fluoro-1,2-benzisoxazol-3-yl)-1-propyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(1H-Indol-3-yl)ethyl)-4-(6-methoxycarbonyl-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-methoxycarbonyl-1H-indol-4-yl)piperazine,  
1-(2-(5-Fluoro-3-benzofuranyl)ethyl)-4-(6-methoxycarbonyl-1H-indol-4-  
30 yl)piperazine,  
1-(5-Fluoro-3-benzofuranylmethyl)-4-(1H-indol-4-yl)piperazine,  
1-(3-Cyano-1H-indol-4-yl)-4-(2-(1H-indol-3-yl)ethyl)piperazine,

- 1-(3-Cyano-1H-indol-4-yl)-4-(2-(5-fluoro-3-benzofuranyl)ethyl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(3-cyano-1H-indol-4-yl)piperazine,  
1-(2-(3-Benzofuranyl)ethyl)-4-(3-cyano-1H-indol-4-yl)piperazine,  
1-(1H-Indol-4-yl)-4-(2-(5-methyl-3-benzofuranyl)ethyl)piperazine,  
5 1-(1H-Indol-4-yl)-4-(2-(4-methyl-3-benzofuranyl)ethyl)piperazine,  
1-(3-(5-Fluoro-3-benzofuranyl)-1-propyl)-4-(1H-indol-4-yl)-1,2,3,6-  
tetrahydropyridine,  
1-(2-(5-Chloro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(1H-Indol-4-yl)-4-(2-(6-methyl-3-benzofuranyl)ethyl)piperazine,  
10 1-(2-(7-Chloro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(4-Chloro-1H-indol-3-yl)ethyl)-4-(3-cyano-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)-4-(1H-indol-4-yl)piperidine,  
1-(2-(5-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(7-Bromo-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
15 1-(2-(4-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(6-Trifluoromethyl-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(1H-Indol-4-yl)-4-(2-(5-methyl-1H-indol-3-yl)ethyl)piperazine,  
1-(1H-Indol-4-yl)-4-(2-(6-methyl-1H-indol-3-yl)ethyl)piperazine,  
1-(1H-Indol-4-yl)-4-(2-(7-methyl-1H-indol-3-yl)ethyl)piperazine,  
20 1-(2-(4,5-Dichloro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(5-Bromo-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(4-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperidine,  
4-(1H-Indol-4-yl)-1-(2-(5-methyl-1H-indol-3-yl)ethyl)piperidine,  
4-(1H-Indol-4-yl)-1-(2-(1H-indol-3-yl)ethyl)piperidine,  
25 1-(1H-Indol-4-yl)-4-(3-(4-methyl-3-benzofuranyl)-1-propyl)piperazine,  
4-(1H-Indol-4-yl)-1-(3-(4-methyl-3-benzofuranyl)-1-propyl)piperidine,  
1-(3-(4-Chloro-3-benzofuranyl)-1-propyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-chloro-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-fluoro-1H-indol-4-yl)piperazine,  
30 1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-cyano-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(7-chloro-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(7-cyano-1H-indol-4-yl)piperazine,

- 1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(2-cyano-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indolin-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-6-yl)piperazine and  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-7-yl)piperazine,  
5 or an acid addition salt thereof.

The invention also relates to a pharmaceutical composition comprising a compound of the invention or a pharmaceutically acceptable acid addition salt thereof and at least one pharmaceutically acceptable carrier or diluent.

10

In another aspect the invention relates to the use of a compound of the invention or a pharmaceutically acceptable acid addition salt thereof for the preparation of a medicament for the treatment of a disorder or disease responsive to the inhibition of serotonin reuptake and antagonism of 5-HT<sub>1A</sub> receptors.

15

- In a final object, the present invention relates to a method for the treatment of a disorder or disease of living animal body, including a human, which is responsive to the inhibition of serotonin reuptake and antagonism of 5-HT<sub>1A</sub> receptors comprising administering to such a living animal body, including a human, a therapeutically  
20 effective amount of a compound as above or a pharmaceutically acceptable acid addition salt thereof.

- Diseases or disorders responsive to the inhibition of serotonin re-uptake and antagonistic activity at 5-HT<sub>1A</sub> receptors include affective disorders, such as  
25 depression, psychosis, anxiety disorders including general anxiety disorder and panic disorder and obsessive compulsive disorder.

- Due to their combined antagonism of 5-HT<sub>1A</sub> receptors and serotonin reuptake inhibiting effect, the compounds of the invention are particularly useful as fast onset  
30 of action medicaments for the treatment of depression. The compounds may also be useful for the treatment of depression in patients who are resistant to treatment with currently available antidepressants.



In the groups of formula (IIA), (IIB) and (IIC), the presence and position of double bonds in the ring containing X, U and Y depend on the meaning of X, U and Y.

Thus, as regards the group of formula (IIA), it is very clear to a person skilled in the art, that when a dotted line emanating from X is a bond, then X is N or CR<sup>1A</sup>, and when the dotted line is not a bond then X is CHR<sup>1A</sup>, NR<sup>1B</sup>, O, or S; and when a dotted line emanating from Y is a bond, then Y is N, or CR<sup>1B</sup> and when the dotted line is not a bond Y is CHR<sup>2A</sup>, NR<sup>2B</sup>, O, or S.

Further, as regards the group of formula (IIB), it is very clear to a person skilled in the art, that when a dotted line emanating from X is a bond then X is N, or CR<sup>1A</sup> and when the dotted line is not a bond then X is CHR<sup>1A</sup>, NR<sup>1B</sup>, O, or S; and when a dotted line emanating from U is a bond then U is C and when the dotted line is not a bond U is CH, or N.

15

And finally as regards the group of formula (IIC), it is very clear to a person skilled in the art that when a dotted line emanating from U is a bond then U is C and when the dotted line is not a bond U is CH, or N; and when a dotted line emanating from Y is a bond then Y is N, or CR<sup>2A</sup> and when the dotted line is not a bond then Y is CHR<sup>2A</sup>, NR<sup>2B</sup>, O, or S.

20

The same applies to W, which is N, CH, or COH when the dotted line emanating from W does not indicate a bond and C when it indicates a bond.

The expression C<sub>1-6</sub>-alk(en/yn)yl means a C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkenyl, or a C<sub>2-6</sub>-alkynyl group. The expression C<sub>3-8</sub>-cycloalk(en)yl means a C<sub>3-8</sub>-cycloalkyl- or cycloalkenyl group.

The term C<sub>1-6</sub> alkyl refers to a branched or unbranched alkyl group having from one to six carbon atoms inclusive, including but not limited to methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl and 2-methyl-1-propyl.

30

Similarly, C<sub>2-6</sub> alkenyl and C<sub>2-6</sub> alkynyl, respectively, designate such groups having from two to six carbon atoms, including one double bond and one triple bond respectively, including but not limited to ethenyl, propenyl, butenyl, ethynyl, propynyl, and butynyl.

5

The terms C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> alkylcarbonyl, hydroxy-C<sub>1-6</sub>-alkyl etc. designate such groups in which the C<sub>1-6</sub> alkyl is as defined above.

- 10 The term C<sub>3-8</sub> cycloalkyl designates a monocyclic or bicyclic carbocycle having three to eight C-atoms, including but not limited to cyclopropyl, cyclopentyl, cyclohexyl, etc.

- The term C<sub>3-8</sub> cycloalkenyl designates a monocyclic or bicyclic carbocycle having  
15 three to eight C-atoms and including one double bond.

In the term C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl and C<sub>1-6</sub>-alk(en/yn)yl are as defined above.

- 20 The term aryl refers to a carbocyclic aromatic group, such as phenyl, naphthyl, in particular phenyl. As used herein aryl may be substituted one or more times with halogen, nitro, cyano, trifluoromethyl, C<sub>1-6</sub>-alkyl, hydroxy and C<sub>1-6</sub>-alkoxy.

- The term heteroaryl refers to a mono- or bicyclic heterocyclic group such as indolyl,  
25 thienyl, pyrimidyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, benzofuranyl, benzothienyl, pyridyl and furanyl, in particular pyrimidyl, indolyl, and thienyl.

- As used herein heteroaryl may be substituted one or more times with halogen, nitro,  
30 cyano, trifluoromethyl, C<sub>1-6</sub>-alkyl, hydroxy and C<sub>1-6</sub>-alkoxy.

In aryl-C<sub>1-6</sub>-alkyl and heteroaryl-C<sub>1-6</sub>-alkyl, aryl, heteroaryl and C<sub>1-6</sub>-alkyl is as defined above.

Halogen means fluoro, chloro, bromo or iodo.

5

As used herein the term acyl refers to formyl, C<sub>1-6</sub>-alk(en/yn)ylcarbonyl, arylcarbonyl, aryl-C<sub>1-6</sub>-alk(en/yn)ylcarbonyl, C<sub>3-8</sub>-cycloalk(en)ylcarbonyl, or a C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl-carbonyl group and the term thioacyl is as the corresponding acyl group in which the carbonyl group is replaced with a thiocarbonyl group.

10

The acid addition salts of the invention are preferably pharmaceutically acceptable salts of the compounds of the invention formed with non-toxic acids. Exemplary of such organic salts are those with maleic, fumaric, benzoic, ascorbic, succinic, oxalic, bis-methylenesalicylic, methanesulfonic, ethanedisulfonic, acetic, propionic, tartaric, 15 salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, and theophylline acetic acids, as well as the 8-halotheophyllines, for example 8-bromotheophylline. Exemplary of such inorganic salts are those with hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, and nitric acids.

20

Further, the compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of this invention.

25

Some of the compounds of the present invention contain chiral centres and such compounds exist in the form of isomers (i.e. enantiomers). The invention includes all such isomers and any mixtures thereof including racemic mixtures.

30

Racemic forms can be resolved into the optical antipodes by known methods, for example, by separation of diastereomeric salts thereof with an optically active acid, and liberating the optically active amine compound by treatment with a base. Another

method for resolving racemates into the optical antipodes is based upon chromatography on an optically active matrix. Racemic compounds of the present invention can also be resolved into their optical antipodes, e.g., by fractional crystallization of d- or l- (tartrates, mandelates, or camphorsulphonate) salts for example. The compounds of the present invention may also be resolved by the formation of diastereomeric derivatives.


Additional methods for the resolution of optical isomers, known to those skilled in the art, may be used. Such methods include those discussed by J. Jaques, A. Collet, and S. Wilen in "Enantiomers, Racemates, and Resolutions", John Wiley and Sons, New York (1981).

Optically active compounds can also be prepared from optically active starting materials.

Finally, formula (I) includes any tautomeric forms of the compounds of the invention.

The compounds of the invention can be prepared by one of the following methods comprising:

- a) reducing the carbonyl groups of a compound of formula (III)




(III)


wherein R<sup>3</sup>-R<sup>12</sup>, W and the dotted lines are as defined above;

19

b) alkylating an amine of formula (IV)



wherein  $R^7$ - $R^{12}$ , W and the dotted lines are as defined above with a reagent of formula (V)




(V)

5


wherein A, Z, n, and m are as defined above and G is a suitable leaving group such as halogen, mesylate, or tosylate;

c) reductive alkylation of an amine of formula

10



wherein  $R^7$ - $R^{12}$ , W and the dotted lines are as defined above with a reagent of formula (VI)




(VI)

15

wherein A, Z, n and m are as defined above and E is either an aldehyde or a carboxylic acid group;


d) reducing the double bond of the indole ring, which is attached to the cyclic amine moiety, of formula (I) in order to obtain the corresponding 2,3-dihydroindole derivatives;

5 e) reducing the double bond of a tetrahydropyridine of formula (VII)



wherein  $R^7$ - $R^{12}$ , A, Z, n, m, and the dotted lines are as previously defined, in order to  
10 obtain the corresponding piperidine derivatives;

f) reducing the amide group of a compound of formula (VIII)




15 wherein  $R^7$ - $R^{12}$ , A, W, Z, n, m, and the dotted lines are as previously defined;

g) reductive removal of one or more of the halogen substituents  $R^3$ - $R^9$  and  $R^{11}$ - $R^{12}$  in a compound of formula (I) in which one or more of these substituents are selected from chloro, bromo, or iodo;


21

h) dialkylating an amine of formula (IX)




wherein  $R^7$ - $R^{12}$  and the dotted line are as defined above with a reagent of formula (X)

5




wherein A, Z, n, and m are as defined above and G is a suitable leaving group such as halogen, mesylate, or tosylate;

10 i) dialkylating an amine of formula (XI)



wherein A, Z, n, and m are as defined above with a reagent of formula (XII)



15

wherein  $R^7$ - $R^{12}$ , W, and the dotted line are as defined above and G is a suitable leaving group such as halogen, mesylate, or tosylate;

- j) alkylating, arylating, or acylating one or both indole nitrogen atoms of a compound of formula (I) in which  $R^{10}$  is hydrogen, and /or X and/or Y is NH; or
- k) reducing a compound of formula (I) in which  $R^7$ ,  $R^8$ , or  $R^9$  is an
- 5 alkoxy carbonyl group in order to obtain the corresponding hydroxymethyl group;

whereupon the compounds of formula (I) are isolated as the free base or in the form of an acid addition salt thereof.

- 10 The reduction according to method a) is preferably carried out in an inert organic solvent such as diethyl ether or tetrahydrofuran in the presence of lithium aluminium hydride at reflux temperature. Starting compounds of formula (III) are generally prepared by condensation of 3-chlorooxalyl indoles (prepared as described in Houben-Weyl, Methoden der Organischen Chemie, Vol E6B2, p. 1058) with amines of
- 15 formula (IV) in the presence of a base such as triethylamine or potassium carbonate.

- The alkylation according to method b) is conveniently performed in an inert organic solvent such as a suitably boiling alcohol or ketone, preferably in the presence of an organic or inorganic base (potassium carbonate or triethylamine) at reflux
- 20 temperature.

- Indolylpiperazine derivatives of formula (IV) are conveniently prepared from the corresponding arylamine according to the method described by Martin et al, J. Med. Chem. 32 (1989) 1052, or the method described by Kruse et al, Rec. Trav. Chim.
- 25 Pays-Bas 107 (1988) 303. The starting arylamines are either commercially available or are well-described in the literature.

- Indolyl tetrahydropyridine derivatives of formula (IV) are known from literature (see eg. French Pat. 2458549). Conveniently, 1-protected 4, 5, 6, or 7-bromoindole is
- 30 lithiated with BuLi followed by addition of 1-protected 4-piperidone and subsequent dehydration as outlined in an example below. The starting bromoindoles are either commercially available or well-described in the literature. Reagents of formula (V) are



either commercially available or can be prepared by literature methods, eg. from the corresponding carboxylic acid derivative by reduction to the corresponding hydroxy derivatives and subsequent conversion of the hydroxy group to the group G by conventional methods.

5

The reductive alkylation according to method c) is performed by standard literature methods. The reaction can be performed in two steps, ie. coupling of derivatives of formula (IV) and the reagent of formula (VI) by standard methods via the carboxylic acid chloride or by use of coupling reagents such as eg. dicyclohexyl carbodiimide followed by reduction of the resulting amide with lithium aluminium hydride or alane. The reaction can also be performed by a standard one-pot procedure. Carboxylic acids or aldehydes of formula (VI) are either commercially available or described in the literature.

15 Reduction of the indole double bond according to method d) is conveniently performed by treatment with diborane or a diborane precursor such as the trimethylamine or dimethylsulfide complex in an inert solvent such as eg. tetrahydrofuran or dioxane from 0 °C to reflux temperature followed by acid catalyzed hydrolysis of the intermediate borane derivative. The reduction can alternatively be performed by treatment with sodium cyanoborohydride in trifluoroacetic acid.

Reduction of the double bonds according to method e) is most conveniently performed by hydrogenation in an alcohol in the presence of a noble metal catalyst, such as eg. platinum or palladium.

25

Reduction of amide groups according to method f) is most conveniently performed with lithium aluminium hydride or alane in an inert organic solvent such as eg. tetrahydrofuran or diethylether from 0 °C to reflux temperature.

30

The removal of halogen substituents according to method g) is conveniently performed by catalytic hydrogenation in an alcohol in the presence of a palladium

catalyst or by treatment with ammonium formate in an alcohol at elevated temperatures in the presence of a palladium catalyst.

The dialkylation of amines according to methods h) and i) is most conveniently performed at elevated temperatures in an inert solvent such as eg. chlorobenzene, toluene, N-methylpyrrolidinone, dimethylformamide, or acetonitrile. The reaction might be performed in the presence of base such as eg. potassium carbonate or triethylamine. Starting material for processes h) and i) are commercially available or can be prepared using conventional methods.

10

The indole N-alkylation or N-acylation are performed in an inert solvent such as eg. an alcohol or ketone at elevated temperatures in the presence of base, eg. potassium carbonate or triethylamine. Alternatively, a phase-transfer reagent can be used. The corresponding N-arylation is best performed under Ullmann-conditions as described in the literature.

15

The reduction of alkoxycarbonyl groups according to method k) is most conveniently performed with lithium aluminium hydride or alane in an inert organic solvent such as e.g. tetrahydrofuran.

20

The following examples will illustrate the invention further. They are, however, not to be construed as limiting.

### Examples

Melting points were determined on a Büchi B-540 apparatus and are uncorrected. Mass spectra were obtained on a Quattro MS-MS system from VG Biotech, Fisons Instruments or on a Sciex API 150EX from Perkin Elmer. Spectra were obtained at two sets of operating conditions using electrospray ionisation: one set to obtain molecular weight information ( $MH^+$ , 20 eV) and the other set to induce fragmentation patterns (70-100 eV). The background was subtracted. The relative intensities of the ions are obtained from the fragmentation pattern. When no intensity is indicated for the molecular ion ( $MH^+$ ) this ion was only present under the first set of operating

30

conditions.  $^1\text{H}$  NMR spectra were recorded at 250 MHz on a Bruker AC 250 or at 500 MHz on a Bruker DRX 500. Deuterated chloroform (99.8% D) or dimethylsulfoxide (99.9% D) were used as solvents. TMS was used as internal reference standard. Chemical shifts are expressed as ppm values. The following abbreviations are used for multiplicity of NMR signals: s=singlet, d=doublet, t=triplet, q=quartet, qv=quintet, h=heptet, dd=double doublet, dt=double triplet, dq=double quartet, tt=triplet of triplets, m= multiplet, b=broad. NMR signals corresponding to acidic protons are to some extent omitted. Content of water in crystalline compounds were determined by Karl Fischer titration. Proper elemental analysis for all target compounds were obtained. Standard work-up procedures refer to extraction with the indicated organic solvent from proper aqueous solutions, drying of combined organic extracts (anhydrous  $\text{MgSO}_4$  or  $\text{NaSO}_4$ ), filtering, and evaporation of the solvent in vacuo. For column chromatography silica gel of type Kieselgel 60, 40-60 mesh ASTM was used.

## 15 Preparation of Intermediates

### A. Preparation of 1-(1H-indol-4-yl)piperazines:

#### 1-(1H-Indol-4-yl)piperazine

Dinitrotoluene (25 g) was dissolved in DMSO (60 mL). Triton-B (40% in methanol, 6.4 mL) was added resulting in a dark purple solution. The mixture was heated to 30 °C followed by slow addition of a solution of paraformaldehyde (4.5 g) in DMSO (40 mL). After addition the mixture was heated to 65 °C for 1.5 hours. Standard work-up with ethyl acetate gave 2-(2,6-dinitrophenyl)ethanol (29 g) as a dark red oil.

The oil (27 g) was dissolved in ethanol (250 mL) and 5% palladium on charcoal (3 g) was added. The mixture was treated with hydrogen gas at 3 atmospheres of pressure in a Parr apparatus for 16 h. Filtration and removal of solvent in vacuo gave 2-(2,6-diaminophenyl)ethanol (19.4 g) as brown oil that crystallized upon standing.

Part of this product (15.8 g) was dissolved in toluen (250 mL) and tris(triphenylphosphine)ruthenium(II)chloride (2.9 g) was added. The mixture was refluxed with a water separator for 8 hours followed by removal of solvent in vacuo.

The remaining reaction mixture was purified by flash chromatography (eluent: heptane/ethyl acetate 3:1) giving 4-amino-1H-indole as a crystalline solid (7.6 g).

The solid was dissolved in chlorobenzene (150 mL) and bis(2-chloroethyl)amine hydrochloride (9 g) was added followed by reflux for 90 hours. Filtration gave a

- 5 crystalline product (9.2 g) which was heated in a mixture of conc. aq. ammonia (50 mL) and ethyl acetate (200 mL) for 15 min. Separation of the organic phase, drying and evaporation gave the title compound as a crystalline material (5.6 g).

The following piperazines were prepared analogously:

- 10 1-(6-Methoxycarbonyl-1H-indol-4-yl)piperazine.  
1-(6-t-Butyl-5-methoxy-1H-indol-4-yl)piperazine.  
1-(6-t-Butyl-7-methoxy-1H-indol-4-yl)piperazine.  
1-(6,7-Dihydro-6,6,8,8-tetramethylcyclopent(g)-1H-indol-4-yl)piperazine

1-(1H-Indol-5-yl)piperazine

- 15 This piperazine was prepared by treatment of 5-aminoindole with and bis(2-chloroethyl)amine by a procedure analogous to the procedure described above for the preparation of 1-(1H-indol-4-yl)piperazine.

1-(3-Cyano-1H-indol-4-yl)piperazine.

- 20 A mixture of 1-(1H-indol-4-yl)piperazine hydrochloride (1 g) and potassium carbonate (2.9 g) in tetrahydrofuran (25 mL) and water (25 mL) was stirred for 15 min followed by addition of a solution of di-t-butyl dicarbonate (2.3 g) in a 1:1 mixture of tetrahydrofuran and water (20 mL). The mixture was stirred at 50 °C for 16 hours. Standard work-up with ethyl acetate gave 1-t-butoxycarbonyl-4-(1H-indol-4-yl)piperazine as a heavy oil (1.1 g).

- 25 The oil was dissolved in acetonitrile (50 mL) followed by dropwise addition of chlorosulfonyl isocyanate (1 mL) at -20 °C. The mixture was kept at low temperature during dropwise addition of dimethylformamide (5 mL) over 20 min. Finally, the mixture was stirred for 30 min at 0 °C. An aqueous solution of sodium carbonate (30 mL) was added and the mixture stirred for 30 min. The organic phase was separated, dried and concentrated. The resulting oil was purified by column chromatography
- 30

(eluent: ethyl acetate/heptane/methanol 16:3:1) giving 1-t-butoxycarbonyl-4-(3-cyano-1H-indol-4-yl)piperazine as a yellow oil (0.5 g).

The oil was dissolved in methanol (2 mL) and ethereal hydrogenchloride (20 mL) was added. Stirring for 2 hours and filtration gave the title compound as a crystalline

5 material (0.34 g).

1-(1-Allyl-1H-indol-4-yl)piperazine.

A solution of 1-t-butoxycarbonyl-4-(1H-indol-4-yl)piperazine (prepared as described above) in tetrahydrofuran (50 mL) was added dropwise to a suspension of sodium  
10 hydride (0.7 g of a 60% mineral oil suspension) in tetrahydrofuran (150 mL) at room temperature. After stirring for further 30 min a solution of allyl bromide (3.5 mL) in tetrahydrofuran (50 mL) was added dropwise. After stirring for 48 h the mixture was poured onto ice-water followed by standard work-up with ethyl acetate giving an oil, which was purified by flash chromatography (eluent: heptane/ethyl acetate 85:15)  
15 giving 1-t-butoxycarbonyl-4-(1-allyl-1H-indol-4-yl)piperazine as an oil (3.2 g).

The oil was dissolved in methanol (15 mL) and a saturated solution of HCl in diethyl ether (100 mL) was added. After stirring for 16 h at room temperature, the resulting colorless crystals consisting of the hydrochloride of the title compound (2.5 g) was collected by filtration and dried in vacuo.

20

The following piperazines were prepared analogously:

1-(1-Benzyl-1H-indol-4-yl)piperazine

1-(1-Propargyl-1H-indol-4-yl)piperazine

25 **B. Preparation of 4-(1H-indol-4-yl)-1,2,3,6-tetrahydropyridine.**

A solution of 4-bromo-1H-indole (36 g) in dimethylformamide (80 mL) was treated with a suspension of NaH (60% in mineraloil, 6.9 g) in dimethylformamide (200 mL) at 20 °C. After stirring for 30 min the mixture was cooled to -10 °C and treated  
30 portionwise with t-butyldimethylsilyl chloride (38 g) followed by stirring for 1 h at room temperature. Standard work-up with ethyl acetate gave an oil which was

purified by flash chromatography giving 4-bromo-1-(t-butyldimethylsilyl)-1H-indole (38 g) as a crystalline material.

The product was dissolved in dry tetrahydrofuran (500 mL), cooled to -78 °C, and treated dropwise with 1.6 M BuLi in hexane (154 mL). After stirring for 30 min at -78 °C, a solution of 1-carbethoxy-4-piperidone (18.2 mL) in tetrahydrofuran (200 mL) was added dropwise followed by stirring for 16 h under slow raise in temperature to room temperature. Standard work-up with diethyl ether gave an oil which was purified by flash chromatography (eluent: heptane/ethyl acetate/triethylamine 6:3:1) giving 1-(t-butyldimethylsilyl)-4-(1-carbethoxy-4-hydroxy-4-piperidinyl)-1H-indole as a crystalline compound (20.5 g).

Treatment of this product with trifluoroacetic acid (15 mL) in methylene chloride (250 mL) at 0 °C for 20-30 min (reaction is followed by thin layer chromatography on silica gel, eluent ethyl acetate/heptane/triethylamine 10:9:1). Addition of 2 M sodium hydroxide, separation of the organic phase, drying, and removal of solvent in vacuo gave an oil, which was purified by flash chromatography (eluent as mentioned above for TLC) giving 1-carbethoxy-4-(1H-indol-4-yl)-1,2,3,6-tetrahydropyridine (9.1 g) as a crystalline material.

Treatment with potassium hydroxide (5 g) in ethanol (150 mL) with a small amount of water (2 mL) at reflux for 3 days gave after standard work-up the title compound as a yellow oil (4.5 g).

### C. Preparation of 5-fluorobenzofuran-3-acetic acid.

A solution of 5-fluorobenzofuran-3-carboxylic acid (56 g) and saturated ethereal solution of hydrochloric gas (300 mL) in methanol (600 mL) was stirred for 16 h at room temperature. Further ethereal HCl was added (300 mL) followed by stirring for 24 h. Concentration in vacuo gave a dark crystalline material, methyl 5-fluorobenzofuran-3-carboxylate (58 g).

Lithium aluminium hydride (15 g) was suspended in tetrahydrofuran (400 mL) under a nitrogen atmosphere followed by dropwise addition of a solution of methyl 5-fluorobenzofuran-3-carboxylate (58 g) in tetrahydrofuran (300 mL). The temperature increased to 55 °C during the addition. After stirring for 2 h the reaction was quenched successively with water (30 mL), 15% aq. sodium hydroxide (15 mL), and

- water (75 mL). Further tetrahydrofuran (500 mL) was added and the mixture stirred for 1 h. The mixture was filtered and the precipitate extracted with a mixture of methylene chloride (1 L) and ethanol (0.5 L). The combined organic phases were concentrated in vacuo giving an oil which was applied to silica gel flash
- 5 chromatography (eluent: methylene chloride/25% aq.  $\text{NH}_3$  99:1). The resulting yellow oil, 5-fluorobenzofuran-3-ylmethanol (14.4 g) crystallised on standing. A solution of 5-fluorobenzofuran-3-ylmethanol (14 g) in methylene chloride (250 mL) was treated succesively with 5 drops of dimethylformamide and thionyl chloride (28 mL). After stirring for 16 h at room temperature the reaction was concentrated in
- 10 vacuo giving 3-chloromethyl-5-fluorobenzofuran as an oil (19.4 g). A suspension of sodium cyanide (10 g) in dimethylsulfoxide (150 mL) was heated to 80 °C followed by quick addition of a solution of 3-chloromethyl-5-fluorobenzofuran (10 g) in dimethylsulfoxide (50 mL). The mixture was kept at 80 °C for ½ h and then poured onto ice. Standard work-up with diethyl ether gave a dark crystalline material,
- 15 5-fluorobenzofuran-3-ylacetonitrile (8.8 g). A solution of 5-fluorobenzofuran-3-ylacetonitrile (8.8 g) in methanol (350 mL) was treated with a saturated ethereal solution of hydrochloric gas (350 mL) and stirred for 16 h at room temperature. The mixture was concentrated in vacuo and standard work-up with diethyl ether/water gave methyl 5-fluorobenzofuran-3-ylacetate (9.4 g) as an
- 20 oil. The obtained methyl ester was dissolved in methanol (200 mL) and 6 M aq. sodium hydroxide (400 mL) was added followed by stirring for 16 at room temperature. Organic solvent was removed in vacuo followed by acidification with 6 M hydrochloric acid. Standard work-up with ethyl acetate gave 5-fluorobenzofuran-3-
- 25 ylacetic acid (9.2 g) as a crystalline material.

The following benzofuran-3-acetic acids were prepared analogously:

2-Methyl-4,5,6,7-tetrafluorobenzofuran-3-acetic acid

Benzofuran-3-acetic acid

30 6-Methylbenzofuran-3-acetic acid

5-Methylbenzofuran-3-acetic acid

4-Methylbenzofuran-3-acetic acid

7-Chlorobenzofuran-3-acetic acid

5-Chlorobenzofuran-3-acetic acid

5-Fluorobenzofuran-3-propionic acid and -butanoic acid, respectively, were prepared

5 by chain prolongation procedures analogously to the procedure described above.

6-Chloroindazol-3-acetic acid was prepared according to J. Med. Chem. 35 (1992) 2155.

3-(6-Fluorobenz[1,2]isoxazol-3-yl)propionic acid was prepared according to Drug Design Discov. 8 (1992) 225.

10

### **Preparation of the compounds of the Invention**

#### **Example 1**

##### **1a, 1-(2-(3-Benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine, oxalate.**

15 A mixture of 2-(3-benzofuranyl)acetic acid (0.95 g), 1-(1H-indol-4-yl)piperazine (1.3 g), and N,N-dicyclohexylcarbodiimide (1.3 g) in a mixture of dry tetrahydrofuran (50 mL) and dry dimethylformamide (10 mL) was stirred for 16 h at room temperature.

Filtration and removal of solvent in vacuo gave an oil which was purified by flash chromatography (eluent: ethyl acetate/heptane/triethylamine 10:9:1) giving 1-(3-

20 benzofuranyl)methylcarbonyl-4-(1H-indol-4-yl)piperazine (0.5 g) as an oil.

The oil was dissolved in tetrahydrofuran (20 mL) and treated with a suspension of lithium aluminium hydride (0.26 g) in tetrahydrofuran (20 mL) at room temperature under a nitrogen atmosphere followed by reflux for 4 hours. The reaction mixture was cooled to 0 °C and treated subsequently with water (1 mL), 15% aq. sodium

25 hydroxide (0.5 mL), and water (2.5 mL). After stirring for 30 min the mixture was

filtered and concentrated. The remaining oil was dissolved in acetone followed by addition of oxalic acid and filtration, giving the title compound as a crystalline

material (0.4 g). Mp 130-32 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.05-3.15 (m, 2H); 3.15-3.30

(m, 6H); 3.35 (s, 4H); 6.45 (s, 1H); 6.55 (d, 1H); 7.00 (t, 1H); 7.10 (d, 1H); 7.20-7.40

30 (m, 3H); 7.60 (d, 1H); 7.75 (d, 1H); 7.90 (s, 1H); 11.10 (s, 1H). MS m/z (%): 346

(MH<sup>+</sup>, 3%), 214 (31%), 199 (19%), 171 (14%).



The following compounds were prepared analogously:

**1b**, *1-(3-Benzofuranylmethyl)-4-(1H-indol-4-yl)piperazine, oxalate*.

Mp 226-28°C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.10-3.20 (m, 4H); 3.20-3.40 (m, 4H); 4.25 (s, 2H); 6.40 (t, 1H); 6.45 (d, 1H); 7.00 (t, 1H); 7.10 (d, 1H); 7.25 (dt, 1H); 7.30-7.45 (m, 2H); 7.65 (dd, 1H); 7.95 (dd, 1H); 8.15 (s, 1H); 11.10 (s, 1H). MS m/z (%): 332 (MH<sup>+</sup>, 10%), 158 (10%), 131 (100%).

**1c**, *1-(2-(5-Fluoro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine, oxalate*.

Mp 196-97 °C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.05 (t, 2H); 3.20-3.45 (m, 10H); 6.44 (s, 1H); 6.50 (d, 1H); 7.00 (t, 1H); 7.10 (d, 1H); 7.20 (dt, 1H); 7.30 (t, 1H); 7.55-7.65 (m, 2H); 8.00 (s, 1H); 11.12 (s, 1H), MS m/z (%) : 364 (MH<sup>+</sup>, 7%), 214 (42%), 199 (20%), 171 (14%).

**1d**, *1-(4-(5-Fluoro-3-benzofuranyl)-1-butyl)-4-(1H-indol-4-yl)piperazine, dihydrochloride*.

Mp 241-44 °C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 1.65-1.95 (m, 4H); 2.70 (t, 2H); 3.15-3.40 (m, 6H); 3.60 (d, 2H); 3.70 (d, 2H); 6.50 (s, 1H); 6.55 (d, 1H); 7.00 (t, 1H); 7.10 (d, 1H); 7.15 (dt, 1H); 7.30 (t, 1H); 7.45-7.60 (m, 2H); 7.90 (s, 1H); 10.95 (b, 1H); 11.20 (s, 1H). MS m/z (%): 392 (MH<sup>+</sup>?, 90%), 234 (19%), 199 (23%), 163 (49%), 131 (11%).

**1e**, *1-(2-(1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine, hemioxalate*.

Mp 167-69 °C, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.05 (s, 4H); 3.15 (s, 4H); 3.30 (s, 4H); 6.40 (s, 1H); 6.50 (d, 1H); 6.90-7.15 (m, 4H); 7.25 (dd, 2H); 7.35 (d, 1H); 7.60 (d, 1H); 10.90 (s, 1H); 11.10 (s, 1H). MS m/z (%): 345 (MH<sup>+</sup>, 5%), 199 (13%), 144 (54%), 107 (9%).

**1f**, *1-(3-(1H-indol-3-yl)-1-propyl)-4-(1H-indol-4-yl)piperazine, oxalate*.

Mp 198-204 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.05 (qv, 2H); 2.75 (t, 2H); 3.15 (t, 2H); 3.35 (s, 8H); 6.45 (s, 1H); 6.50 (d, 1H); 7.00 (t, 2H); 7.10 (t, 2H); 7.20 (d, 1H); 7.25 (t, 1H); 7.35 (d, 1H); 7.55 (d, 1H); 10.85 (b, 1H); 11.15 (b, 1H).

**1g**, *1-(4-(1H-indol-3-yl)-1-butyl)-4-(1H-indol-4-yl)piperazine, oxalate*.

Mp 189-93 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 1.60-1.85 (m, 4H); 2.75 (t, 2H); 3.05 (t, 2H); 3.15-3.50 (m, 8H); 6.45 (s, 1H); 6.50 (s, 1H); 6.90-7.20 (m, 5H); 7.30 (s, 1H); 7.35 (d, 1H); 7.55 (d, 1H); 8.20 (b, 2H); 10.90 (s, 1H); 11.20 (s, 1H).

5

**1h**, *1-(3-(5-Fluoro-3-benzofuranyl)-1-propyl)-4-(1H-indol-4-yl)piperazine, dihydrochloride*.

Mp 230-34 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.20 (qv, 2H); 2.75 (t, 2H); 3.20-3.30 (m, 2H); 3.30-3.45 (m, 4H); 3.55-3.80 (m, 4H); 6.55 (s, 1H); 6.65 (d, 1H); 7.00 (t, 1H); 7.10-7.20 (m, 2H); 7.30 (t, 1H); 7.50-7.65 (m, 2H); 7.95 (s, 1H); 11.25 (s, 1H); 11.40 (b, 1H). MS m/z (%): 378 (MH<sup>+</sup>, 70%), 220 (26%), 199 (25%), 177 (18%), 159 (100%).

10

**1i**, *1-(2-(2-Methyl-4,5,6,7-tetrafluoro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine, dihydrochloride*.

Mp 181-87 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.50 (s, 3H); 3.15-3.50 (m, 8H); 3.65-3.85 (m, 4H); 6.50 (s, 1H); 6.60 (t, 1H); 7.05 (t, 1H); 7.15 (d, 1H); 7.35 (d, 1H); 11.20 (s, 1H); 11.75 (b, 1H). MS m/z (%): 414 (MH<sup>+</sup> - F, 13%), 396 (11%), 214 (72%), 199 (23%), 195 (34%), 159 (26%).

20

**1j**, *1-(2-(3-Indazolyl)ethyl)-4-(1H-indol-4-yl)piperazine, oxalate*.

Mp 149-51 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.05-3.15 (m, 4H); 3.15-3.20 (m, 2H); 3.20-3.35 (m, 6H); 6.40 (s, 1H); 6.50 (d, 1H); 7.00 (t, 1H); 7.05 (d, 1H); 7.10 (t, 1H); 7.25 (s, 1H); 7.35 (t, 1H); 7.50 (d, 1H); 7.80 (d, 1H); 11.10 (s, 1H). MS m/z (%): 346 (MH<sup>+</sup>, 40%); 199 (80%); 144 (100%).

25

**1k**, *1-(2-(6-Chloro-3-indazolyl)ethyl)-4-(1H-indol-4-yl)piperazine, oxalate*.

Mp 255-58 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.10-3.15 (m, 4H); 3.15-3.20 (m, 2H); 3.20-3.30 (m, 6H); 6.40 (s, 1H); 6.50 (d, 1H); 7.00 (t, 1H); 7.10 (d, 1H); 7.15 (d, 1H); 7.25 (s, 1H); 7.55 (s, 1H); 7.85 (d, 1H); . MS m/z (%): 380 (MH<sup>+</sup>, 100%), 214 (30%), 139 (50%).

30

**11**, *1-(2-(7-Cyano-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine, oxalate*.

Mp 241-43 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.15 (t, 2H); 3.30 (t, 2H); 3.30-3.50 (m, 8H); 6.45 (s, 1H); 6.50 (d, 1H); 7.00 (t, 1H); 7.10 (d, 1H); 7.20 (t, 1H); 7.30 (s, 1H); 7.50 (s, 1H); 7.60 (d, 1H); 8.00 (d, 1H); 11.15 (s, 1H); 11.90 (s, 1H). MS m/z (%): 370 (MH<sup>+</sup>, 100%), 214 (28%), 156 (42%).

## Example 2

**2a**, *1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine, oxalate*.

10 A solution of 6-chloro-1H-indole (15 g) in diethyl ether (300 mL) was cooled to 0 °C and treated with a solution of oxalyl chloride (9.4 mL) in diethyl ether (30 mL). The mixture was stirred for 16 hours at room temperature. Filtration gave 2-(6-chloro-1H-indol-3-yl)-2-oxoacetyl chloride as a crystalline material (15.5 g).

A part of this product (2.5 g) was dissolved in dry tetrahydrofuran (25 mL) and added dropwise to a solution of 1-(1H-indol-4-yl)piperazine (1.4 g) and triethylamine (15 mL) in tetrahydrofuran (100 mL). After stirring for 16 hours the reaction mixture was concentrated in vacuo. The remaining oil was purified by flash chromatography (eluent: ethyl acetate/methanol/triethylamine 85:10:5) giving 1-(2-(6-chloro-1H-indol-3-yl)-1,2-dioxoethyl)-4-(1H-indol-4-yl)piperazine (1.6 g) as a crystalline material.

20 This product was suspended in tetrahydrofuran (25 mL) and added dropwise to a suspension of lithium aluminium hydride (1.5 g) in tetrahydrofuran (50 mL). The mixture was refluxed for 4 hours and cooled to 0 °C followed by subsequent addition of water (3 mL), 15% aq. sodium hydroxide (1.5 mL), and water (7.5 mL). Filtration and standard work-up gave a yellow oil which was converted to the title oxalate salt (1.5 g) from an acetone solution by addition of oxalic acid. Mp 229-31 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.10 (t, 2H); 3.25-3.55 (m, 10H); 6.45 (s, 1H); 6.50 (d, 1H); 6.90-7.10 (m, 3H); 7.25-7.35 (m, 2H); 7.45 (s, 1H); 7.65 (d, 1H); 11.12 (s, 2H), MS m/z (%): 379 (MH<sup>+</sup>, 18%), 214 (16%), 199 (17%), 178 (16%), 143 (13%).

30 The following compounds were prepared analogously:

**2b**, 1-(2-(4-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine, oxalate.

Mp 245-47 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.30-3.50 (m, 12H); 6.45 (s, 1H); 6.50 (d, 1H); 6.95-7.10 (m, 4H); 7.30 (s, 1H); 7.35-7.40 (m, 2H); 11.15 (s, 1H); 11.40 (s, 1H). MS m/z (%): 379 (MH<sup>+</sup>, 28%), 178 (41%), 143 (100%).

5

**2c**, 1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine, fumarate.

Mp 212-15 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.75-3.10 (m, 8H); 3.10-3.35 (m, 4H); 6.40 (s, 1H); 6.50 (d, 1H); 6.60 (s, 2H); 6.85-7.10 (m, 3H); 7.20-7.40 (m, 4H); 10.95 (s, 1H); 11.05 (s, 1H). MS m/z (%): 363 (MH<sup>+</sup>, 18%), 214 (100%), 202 (34%), 199 (19%), 171 (12%), 162 (87%).

10

**2g**, 1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)-1,2,3,6-tetrahydropyridine, 1.5 fumarate.

Mp 225-26 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.60-2.70 (m, 2H); 2.85-3.10 (m, 6H); 3.40-3.50 (m, 2H); 6.10 (s, 1H); 6.60 (s, 3H); 6.90-7.10 (m, 3H); 7.20-7.40 (m, 3H); 7.40 (d, 1H); 7.60 (d, 1H); 11.05 (s, 1H); 11.15 (s, 1H). MS m/z (%): 376 (MH<sup>+</sup>, 12%), 179 (80%), 143 (100%).

15

**2h**, 1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)-1,2,3,6-tetrahydropyridine, trifumarate.

Mp 204-6 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.70-2.85 (m, 2H); 3.00-3.20 (m, 4H); 3.25 (t, 2H); 3.70-3.80 (m, 2H); 6.05-6.15 (m, 1H); 6.60 (s, 6H); 6.85-7.00 (m, 2H); 7.10 (t, 1H); 7.30-7.45 (m, 6H); 11.00 (s, 1H); 11.20 (s, 1H). MS m/z (%): 360 (MH<sup>+</sup>), 162 (100%).

25

**2i**, 1-(2-(7-Bromo-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine, hemioxalate.

Mp 149-51 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.95-3.20 (m, 8H); 3.20-3.40 (m, 4H); 6.40 (s, 1H); 6.50 (d, 1H); 7.00 (dt, 2H); 7.10 (d, 1H); 7.25 (t, 1H); 7.30-7.40 (m, 2H); 7.60 (d, 1H); 11.05 (s, 1H); 11.10 (s, 1H). MS m/z (%): 423 (MH<sup>+</sup>, 9%), 222 (20%), 214 (29%), 143 (100%).

30

**2j**, 1-(1-Allyl-1H-indol-4-yl)-4-(2-(6-chloro-1H-indol-3-yl)ethyl)piperazine, fumarate.

Mp 230-32 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.70-3.00 (m, 8H); 3.10-3.30 (m, 4H); 4.75 (d, 2H); 5.00 (d, 1H); 5.15 (d, 1H); 5.85-6.05 (m, 1H); 6.40 (d, 1H); 6.55 (dd, 1H); 6.60 (s, 2H); 6.95-7.10 (m, 3H); 7.20-7.30 (m, 2H); 7.40 (d, 1H); 7.55 (d, 1H). MS m/z

5 (%) : 419 (MH<sup>+</sup>, 13%), 254 (21%), 143 (100%).

**2k**, 1-(1-Allyl-1H-indol-4-yl)-4-(2-(5-fluoro-1H-indol-3-yl)ethyl)piperazine, 1.25

fumarate.

Mp 210-12 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.75-3.00 (m, 8H); 3.10-3.30 (m, 4H); 4.80 (d, 2H); 5.00 (d, 1H); 5.15 (d, 1H); 5.90-6.10 (m, 1H); 6.40 (d, 1H); 6.50 (dd, 1H); 6.60 (s, 2H); 6.90 (dt, 1H); 7.00-7.10 (m, 1H); 7.20-7.40 (m, 4H); 10.95 (s, 1H). MS m/z

10 (%) : 403 (MH<sup>+</sup>, 25%), 239 (30%), 162 (100%).

**2l**, 1-(1-Benzyl-1H-indol-4-yl)-4-(2-(6-chloro-1H-indol-3-yl)ethyl)piperazine,

15 hemifumarate.

Mp 237-39 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.65-2.85 (m, 6H); 2.90 (t, 2H); 3.10-3.25 (m, 4H); 5.40 (s, 2H); 6.45 (d, 1H); 6.50 (d, 1H); 6.60 (s, 1H); 6.95-7.10 (m, 3H); 7.15 (d, 1H); 7.20-7.35 (m, 5H); 7.35-7.45 (m, 2H); 7.60 (d, 1H); 11.00 (s, 1H). MS m/z (%):

469 (MH<sup>+</sup>, 20%), 304 (32%), 289 (22%), 143 (100%).

20

**2m**, 1-(1-Benzyl-1H-indol-4-yl)-4-(2-(5-fluoro-1H-indol-3-yl)ethyl)piperazine,

fumarate.

Mp 178-80 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.70-3.00 (m, 8H); 3.10-3.30 (m, 4H); 5.40 (s, 2H); 6.45 (d, 1H); 6.50 (d, 1H); 6.60 (s, 2H); 6.90 (dt, 1H); 7.00 (d, 1H); 7.10 (d, 1H);

25 7.15 (d, 1H); 7.20-7.40 (m, 5H); 7.45 (d, 1H); 10.95 (s, 1H). MS m/z (%): 453 (MH<sup>+</sup>, 28%), 304 (39%), 162 (100%).

**2n**, 1-(1-Benzyl-1H-indol-4-yl)-4-(2-(5-bromo-1H-indol-3-yl)ethyl)piperazine,

fumarate.

30 Mp 230-32 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.75-3.05 (m, 8H); 3.10-3.35 (m, 4H); 5.45 (s, 2H); 6.45 (d, 1H); 6.50 (d, 1H); 6.60 (s, 2H); 7.00 (t, 1H); 7.10 (d, 1H); 7.10-7.20 (m,

2H); 7.20-7.40 (m, 5H); 7.40 (d, 1H); 7.80 (s, 1H); 11.05 (s, 1H). MS m/z (%): 513 (MH<sup>+</sup>, 14%), 304 (30%), 142 (100%).

**2o**, *1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine*.

- 5 **Mp** 197-99 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.70-2.95 (m, 6H); 3.00 (t, 2H); 3.20-3.40 (m, 5H); 4.85 (d, 2H); 6.50 (d, 1H); 6.65 (d, 1H); 7.00-7.30 (m, 5H); 7.35 (s, 1H); 7.55 (d, 1H); 8.00 (s, 1H). MS m/z (%): 417 (MH<sup>+</sup>, 15%), 252 (24%), 237 (17%), 143 (100%).

- 10 **2p**, *1-(2-(1H-Indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine, hemifumarate*.

- Mp** 193-95 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.60-2.85 (m, 6H); 2.90 (t, 2H); 3.10-3.25 (m, 4H); 3.35 (t, 1H); 5.00 (d, 2H); 6.45 (d, 1H); 6.55 (d, 1H); 6.60 (s, 1H); 6.90-7.25 (m, 5H); 7.25-7.40 (m, 2H); 7.55 (d, 1H), 10.75 (s, 1H). MS m/z (%): 383 (MH<sup>+</sup>, 44%),  
15 252 (55%), 143 (100%).

**2q**, *1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine*.

- Mp** 153-55 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.70-2.90 (m, 6H), 2.90-3.10 (m, 3H); 3.25-3.45 (m, 4H); 4.85 (d, 2H); 6.55 (d, 1H); 6.65 (d, 1H); 6.95 (dt, 1H); 7.00-7.35 (m, 6H);  
20 8.00 (s, 1H). MS m/z (%): 401 (MH<sup>+</sup>, 48%), 237 (27%), 162 (81%), 115 (100%).

**2r**, *1-(2-(5-Bromo-1H-indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine*.

- Mp** 154-56 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.70-2.90 (m, 6H); 2.90-3.00 (m, 3H); 3.25-3.40 (m, 4H); 4.85 (d, 2H); 6.55 (d, 1H); 6.65 (d, 1H); 7.00-7.10 (m, 2H); 7.10-7.35 (m,  
25 4H); 7.75 (s, 1H); 8.05 (s, 1H). MS m/z (%): 461 (MH<sup>+</sup>, 5%), 252 (16%), 237 (12%), 143 (100%).

**2s**, *1-(1-Benzyl-1H-indol-4-yl)-4-(2-(1H-indol-3-yl)ethyl)piperazine, hemifumarate*.

- Mp** 188-90 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.65-2.85 (m, 6H); 2.95 (t, 2H); 3.10-3.30 (m,  
30 4H); 5.45 (s, 2H); 6.45 (d, 1H); 6.50 (d, 1H); 6.60 (s, 1H); 6.90-7.10 (m, 4H); 7.10-7.35 (m, 7H); 7.40 (d, 1H); 7.55 (d, 1H); 10.80 (s, 1H). MS m/z (%): 435 (MH<sup>+</sup>, 22%), 304 (52%), 143 (100%).

**2t**, 1-(2-(5-Bromo-1H-indol-3-yl)ethyl)-4-(1H-indol-5-yl)piperazine, fumarate.

Mp 231-33 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.80 (t, 2H); 2.80-2.90 (m, 4H); 2.95 (t, 2H); 3.10-3.20 (m, 4H); 6.30 (s, 1H); 6.60 (s, 2H); 6.85 (dd, 1H); 7.05 (s, 1H); 7.20 (dd, 1H); 7.25 (t, 1H); 7.25-7.30 (m, 2H); 7.35 (d, 1H); 7.75 (s, 1H); 10.80 (s, 1H); 11.05 (s, 1H). MS m/z (%): 425 (MH<sup>+</sup>, 11%), 223 (14%), 143 (100%).

**2u**, 1-(2-(5-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-5-yl)piperazine, hemifumarate.

Mp 232-34 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.70 (t, 2H); 2.70-2.80 (m, 4H); 2.90 (t, 2H); 3.05-3.15 (m, 4H); 6.30 (s, 1H); 6.60 (s, 1H); 6.85 (dd, 1H); 7.00 (s, 1H); 7.05 (d, 1H); 7.25 (t, 1H); 7.25-7.30 (m, 2H); 7.35 (d, 1H); 7.60 (s, 1H); 10.80 (s, 1H); 11.00 (s, 1H). MS m/z (%): 379 (MH<sup>+</sup>, 18%), 143 (100%).

### Example 3

**3a**, 1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(6-hydroxymethyl-1H-indol-4-yl)piperazine.

A solution of 1-(2-(5-fluoro-1H-indol-3-yl)-1,2-dioxoethyl)-4-(6-methoxycarbonyl-1H-indol-4-yl)piperazine (1.8 g, prepared from 2-(5-fluoro-1H-indol-3-yl)-2-oxoacetyl chloride and 1-(6-methoxycarbonyl-1H-indol-4-yl)piperazine by the procedure described in Example 2) in tetrahydrofuran (50 mL) was added dropwise to a suspension of lithium aluminium hydride (1.7 g) in tetrahydrofuran (125 mL) at room temperature followed by reflux for 4 hours. The reaction mixture was cooled to 5 °C followed by subsequent addition of water (3.4 mL), 15% aq. sodium hydroxide (1.7 mL), and water (8.5 mL). Filtration and removal of solvent in vacuo gave an oil which was purified by flash chromatography (eluent: ethyl acetate/methanol/triethylamine 85:10:5) giving the title product (0.9 g), which was crystallized from diisopropyl ether. Mp 198-200 °C. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 2.60-2.80 (m, 6H); 2.85 (t, 2H); 3.15 (s, 4H); 4.45-4.55 (m, 2H); 4.90-5.00 (m, 1H); 6.30 (s, 1H); 6.40 (s, 1H); 6.90 (dd, 1H); 7.00 (s, 1H); 7.20 (s, 1H); 7.25-7.35 (m, 3H); 10.85 (s, 1H); 10.95 (s, 1H).

The following compounds were prepared analogously:

**3b**, *1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-hydroxymethyl-1H-indol-4-yl)piperazine*.

Mp 194-96 °C. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 2.60-2.80 (m, 6H); 2.90 (t, 2H); 3.1-3.20 (m, 4H); 4.50 (d, 2H), 4.95 (t, 1H), 6.35 (s, 1H); 6.45 (s, 1H); 6.95-7.05 (m, 1H); 7.20 (t, 1H), 7.25 (d, 1H), 7.35 (d, 1H), 7.55 (d, 1H); 10.95 (s, 2H).

**3c**, *1-(2-(5-Bromo-1H-indol-3-yl)ethyl)-4-(6-hydroxymethyl-1H-indol-4-yl)piperazine*.

Mp 163-65 °C. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 2.65 (t, 2H); 2.70 (s, 4H); 2.90 (t, 2H); 3.15 (s, 4H); 4.50 (d, 2H); 4.95 (t, 1H); 6.35 (s, 1H); 6.45 (s, 1H); 7.00 (s, 1H); 7.10-7.20 (m, 2 H); 7.25 (s, 1H); 7.30 (d, 1H); 7.70 (s, 1H); 10.90 (s, 1H); 11.00 (s, 1H).

#### Example 4

**4a**, *1-(3-(6-Fluoro-1,2-benzisoxazol-3-yl)-1-propyl)-4-(1H-indol-4-yl)piperazine fumarate*.

A solution of 3-(3-bromo-1-propyl)-6-fluoro-1,2-benzisoxazole (1.1 g), 1-(1H-indol-4-yl)piperazine (1.0 g), potassium carbonate (1.9 g), and potassium iodide (50 mg) in 4-methyl-2-pentanone (100 mL) was refluxed for 16 hours. Filtration and removal of solvent in vacuo gave an oil which was purified by flash chromatography (eluent: heptane/ethyl acetate/triethylamine 75:20:5) giving an oil (1.0 g) which was crystallized as the title fumarate from acetone by addition of fumaric acid. Mp 187-89 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.00 (qv, 2H); 2.55 (t, 2H); 2.60-2.80 (m, 4H); 3.05 (t, 2H); 3.05-3.20 (m, 4H); 6.40 (s, 1H); 6.45 (d, 1H); 6.60 (s, 2H); 6.90-7.05 (m, 2H); 7.20-7.35 (m, 2H); 7.70 (dd, 1H); 8.00 (dd, 1H); 11.00 (s, 1H). MS m/z (%): 379 (MH<sup>+</sup>, 10%), 178 (100%), 159 (24%).

The following compounds were prepared analogously:

**4f**, *1-(2-(1H-Indol-3-yl)ethyl)-4-(6-methoxycarbonyl-1H-indol-4-yl)piperazine oxalate*.



Mp 213-16 °C. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 3.10 (t, 2H); 3.20-3.60 (m, 10H); 3.80 (s, 3H); 6.55 (s, 1H); 6.90-7.10 (m, 3H); 7.25 (s, 1H); 7.35 (d, 1H); 7.50 (s, 1H); 7.60 (d, 1H); 7.75 (s, 1H); 10.90 (s, 1H); 11.55 (s, 1H).

- 5 **4g**, 1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-methoxycarbonyl-1H-indol-4-yl)piperazine, oxalate.

Mp 228-30 °C. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 3.10 (t, 2H); 3.30 (t, 2H); 3.45 (s, 8H); 3.85 (s, 3H); 6.55 (s, 1H); 7.05 (dd, 1H); 7.10 (s, 1H); 7.33 (s, 1H); 7.45 (s, 1H); 7.55 (t, 1H); 7.65 (d, 1H); 7.85 (s, 1H); 11.15 (s, 1H); 11.65 (s, 1H).

10

- 4h**, 1-(2-(5-Fluoro-3-benzofuranyl)ethyl)-4-(6-methoxycarbonyl-1H-indol-4-yl)piperazine, oxalate.

Mp 227-28 °C. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 3.05 (t, 2H); 3.25 (t, 2H); 3.25-3.35 (m, 4H); 3.35-3.45 (m, 4H); 3.85 (s, 3H); 6.55 (s, 1H); 7.10 (s, 1H); 7.15 (t, 1H); 7.55 (t, 1H);

- 15 7.55-7.65 (m, 2H); 7.80 (s, 1H); 8.00 (s, 1H); 11.55 (s, 1H).

- 4i**, 1-(5-Fluoro-3-benzofuranylmethyl)-4-(1H-indol-4-yl)piperazine, dihydrochloride.

Mp 238-40 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.20-3.50 (m, 4H); 3.60 (d, 2H); 3.75 (d, 2H); 4.60 (s, 2H); 6.50 (s, 1H); 6.55 (d, 1H); 7.00 (t, 1H); 7.15 (d, 1H); 7.25 (dt, 1H); 7.25-  
20 7.30 (m, 1H); 7.70 (dd, 1H); 8.00 (dd, 1H); 8.40 (s, 1H); 11.22 (s, 1H); 11.65 (b, 1H).  
MS m/z (%): 350 (MH<sup>+</sup>, 7%), 201 (34%), 159 (100%), 149 (20%).

- 4m**, 1-(3-Cyano-1H-indol-4-yl)-4-(2-(1H-indol-3-yl)ethyl)piperazine, fumarate.

Mp >250 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.70-3.00 (m, 8H); 3.10 (s, 4H); 6.60 (s, 1H); 6.75  
25 (dd, 1H); 6.95-7.10 (m, 2H); 7.15-7.20 (m, 3H); 7.35 (d, 1H); 8.20 (s, 1H); 10.80 (s,  
1H); 12.20 (s, 1H). MS m/z (%): 370 (MH<sup>+</sup>, 9%), 239 (100%), 227 (25%), 224  
(27%), 144 (65%).

- 4n**, 1-(3-Cyano-1H-indol-4-yl)-4-(2-(5-fluoro-3-benzofuranyl)ethyl)piperazine,  
30 hemifumarate.

Mp 235-37 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.60-2.95 (m, 8H); 3.10 (s, 4H); 6.60 (s, 2H);  
6.75 (d, 1H); 7.05-7.25 (m, 3H); 7.45-7.60 (m, 2H); 7.95 (s, 1H); 8.20 (s, 1H); 12.15

(s, 1H). MS m/z (%): 389 (MH<sup>+</sup>, 8%), 239 (100%), 224 (38%), 208 (14%), 163 (15%).

**4o**, 1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(3-cyano-1H-indol-4-yl)piperazine,

5 *hemifumarate*.

Mp 234-36 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.70 (t, 2H); 2.80 (b, 4H); 2.90 (t, 2H); 3.10 (b, 4H); 6.60 (d, 2H); 6.75 (d, 1H); 7.00 (d, 1H); 7.15-7.20 (m, 2H); 7.25 (s, 1H); 7.35 (s, 1H); 7.55 (d, 1H); 8.20 (s, 1H); 10.90 (s, 1H); 12.10 (s, 1H). MS m/z (%): 404 (MH<sup>+</sup>, 8%), 239 (100%), 227 (18%), 224 (30%), 178 (25%).

10

**4p**, 1-(2-(3-Benzofuranyl)ethyl)-4-(3-cyano-1H-indol-4-yl)piperazine, *sesquioxalate*.

Mp 221-23 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.10 (t, 2H); 3.20-3.45 (m, 10 H); 6.80 (d, 1H); 7.15-7.40 (m, 4H); 7.60 (dd, 1H); 7.75 (dd, 1H); 7.90 (s, 1H); 8.25 (d, 1H); 12.30 (s, 1H). MS m/z (%): 371 (MH<sup>+</sup>, 20), 239 (63%), 145 (100%).

15

**4q**, 1-(1H-Indol-4-yl)-4-(2-(5-methyl-3-benzofuranyl)ethyl)piperazine, *hydrochloride*.

Mp 258-60 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.40 (s, 3H); 3.15-3.55 (m, 8H); 3.65-3.80 (m, 4H); 6.50 (d, 1H); 6.60 (d, 1H); 7.00 (t, 1H); 7.15 (d, 1H); 7.20 (d, 1H); 7.30 (t, 1H); 7.45 (d, 1H); 7.60 (s, 1H); 7.90 (s, 1H); 11.20 (s, 1H). MS m/z (%): 360 (MH<sup>+</sup>, 10%),

20 214 (97%), 143 (100%).

**4r**, 1-(1H-Indol-4-yl)-4-(2-(4-methyl-3-benzofuranyl)ethyl)piperazine, *oxalate*.

Mp 204-6 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.05 (s, 3H); 3.30-3.50 (m, 12H); 6.45 (d, 1H); 6.50 (d, 1H); 6.95-7.05 (m, 2H); 7.10 (d, 1H); 7.20 (t, 1H); 7.30 (t, 1H); 7.40 (d, 1H);  
25 7.85 (s, 1H); 11.10 (s, 1H). MS m/z (%): 360 (MH<sup>+</sup>, 23%), 214 (81%), 199 (100%), 143 (53%).

**4s**, 1-(3-(5-Fluoro-3-benzofuranyl)-1-propyl)-4-(1H-indol-4-yl)-1,2,3,6-tetrahydropyridine, *fumarate*.

30 Mp 183-85 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 1.70-2.00 (m, 2H); 2.40-2.90 (m, 8H); 3.20-3.35 (m, 2H); 6.00-6.10 (m, 1H); 6.55 (s, 1H); 6.60 (s, 2H); 6.90 (d, 1H); 7.05 (t, 1H); 7.15

(dt, 1H); 7.25-7.40 (m, 2H); 7.50 (dd, 1H); 7.55 (dd, 1H); 7.90 (s, 1H); 11.10 (s, 1H).  
MS m/z (%): 375 (MH<sup>+</sup>, 90%), 206 (100%), 149 (90%).

**4t**, *1-(2-(5-Chloro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine, oxalate*.

- 5   Mp 200-2 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 3.10 (t, 2H); 3.30-3.50 (m, 10H); 6.45 (d, 1H);  
6.50 (d, 1H); 7.00 (t, 1H); 7.10 (d, 1H); 7.30 (t, 1H); 7.40 (dd, 1H); 7.65 (d, 1H); 7.90  
(s, 1H); 8.00 (s, 1H); 11.10 (s, 1H). MS m/z (%): 380 (MH<sup>+</sup>, 50%), 214 (100%), 143  
(80%).

- 10   **4u**, *1-(1H-Indol-4-yl)-4-(2-(6-methyl-3-benzofuranyl)ethyl)piperazine, oxalate*.

    Mp 190-92 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.10 (s, 3H); 3.10 (t, 2H); 3.25-3.50 (m, 10H);  
6.45 (d, 1H); 6.50 (d, 1H); 7.00 (t, 1H); 7.10 (t, 2H); 7.30 (t, 1H); 7.40 (s, 1H); 7.65  
(d, 1H); 7.80 (s, 1H); 11.15 (s, 1H). MS m/z (%): 360 (MH<sup>+</sup>, 14%), 214 (44%), 143  
(100%).

15

**4v**, *1-(2-(7-Chloro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine, hemioxalate*.

- Mp 216-18 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.85-3.15 (m, 8H); 3.15-3.40 (m, 4H); 6.40 (d,  
1H); 6.50 (d, 1H); 7.00 (t, 1H); 7.05 (d, 1H); 7.25 (t, 1H); 7.30 (d, 1H); 7.45 (d, 1H);  
7.75 (d, 1H); 8.00 (s, 1H); 11.05 (s, 1H). MS m/z (%): 380 (MH<sup>+</sup>, 13%), 199 (30%),  
20   143 (100%).

**4x**, *1-(2-(4-Chloro-1H-indol-3-yl)ethyl)-4-(3-cyano-1H-indol-4-yl)piperazine,  
hemifumarate*.

- Mp >250 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 2.65-3.00 (m, 6H); 3.00-3.30 (m, 6H); 6.60 (s,  
25   1H); 6.75 (dd, 1H); 6.90-7.10 (m, 2H); 7.10-7.25 (m, 2H); 7.25-7.40 (m, 2H); 8.20 (d,  
1H); 11.20 (s, 1H); 12.10 (s, 1H). MS m/z (%): 404 (MH<sup>+</sup>, 14%), 224 (15%), 184  
(18%), 143 (100%).

**Example 5****5a, 1-(2-(6-Chloro-1H-indol-3-yl)-4-(1H-indol-4-yl)piperidine.**

A solution of compound **2g** (0.5 g) was dissolved in glacial acetic acid. Platinum catalyst (10% on charcoal, 10 mg) was added and the mixture hydrogenated in a Parr apparatus under 3 atmospheres of hydrogen gas for 16 hours. Filtration and removal of solvent gave the title compound as a crystalline material (0.4 g). Mp 205-6 °C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 1.75-1.90 (m, 4H); 2.10-2.20 (m, 2H); 2.60-2.70 (m, 2H); 2.85-2.95 (m, 3H); 3.15 (d, 2H); 6.50 (s, 1H); 6.80-6.90 (m, 2H); 6.95-7.10 (m, 2H); 7.20 (s, 1H); 7.30 (s, 1H); 7.40 (s, 1H); 7.55 (d, 1H); 10.90 (s, 1H); 11.00 (s, 1H). MS m/z (%): 378 (MH<sup>+</sup>, 18%), 178 (100%), 143 (47%).

**Pharmacological Testing**

- 15 The affinity of the compounds of the invention to 5-HT<sub>1A</sub> receptors was determined by measuring the inhibition of binding of a radioactive ligand at 5-HT<sub>1A</sub> receptors as described in the following test:

**Inhibition of <sup>3</sup>H-5-CT Binding to Human 5-HT<sub>1A</sub> Receptors.**

20

- By this method the inhibition by drugs of the binding of the 5-HT<sub>1A</sub> agonist <sup>3</sup>H-5-carboxamido tryptamine (<sup>3</sup>H-5-CT) to cloned human 5-HT<sub>1A</sub> receptors stably expressed in transfected HeLa cells (HA7) (Fargin, A. *et al*, *J. Biol. Chem.*, **1989**, 264, 14848) is determined *in vitro*. The assay was performed as a modification of the method described by Harrington, M.A. *et al*, *J. Pharmacol. Exp. Ther.*, **1994**, 268, 1098. Human 5-HT<sub>1A</sub> receptors (40 µg of cell homogenate) were incubated for 15 minutes at 37 °C in 50 mM Tris buffer at pH 7.7 in the presence of <sup>3</sup>H-5-CT. Non-specific binding was determined by including 10 µM of metergoline. The reaction was terminated by rapid filtration through Unifilter GF/B filters on a Tomtec Cell Harvester. Filters were counted in a Packard Top Counter. The results obtained are presented in table 1:
- 30

Compound No.	Inhibition of $^3\text{H}$ -5-CT Binding $\text{IC}_{50}$ (nM)	Compound No.	Inhibition of $^3\text{H}$ -5-CT Binding $\text{IC}_{50}$ (nM)
1a	10	3b	120
1b	210	3c	1000
1c	12	4a	9.8
1d	2.5	4f	1200
1e	6.9	4g	350
1f	9.8	4h	3100
1g	13	4l	330
1h	11	4m	3.2
1i	3.4	4n	5.7
1j	22	4o	2.9
1k	22	4p	4.3
1l	2.5	4q	21
2a	3.4	4r	10
2b	11	4s	16
2c	27	4t	9.4
2g	2.8	4u	7.0
2h	11	4v	4.3
2i	5.4	4x	5.4
2t	230	5a	14
2u	310		
3a	910	Pindolol*	100

Table 1 \* reference compound

The compounds of the invention have also been tested for their effect on re-uptake of

5 serotonin in the following test:

**Inhibition of  $^3\text{H}$ -5-HT Uptake Into Rat Brain Synaptosomes.**

Using this method the ability of drugs to inhibit the accumulation of  $^3\text{H}$ -5-HT into whole rat brain synaptosomes is determined *in vitro*. The assay was performed as

5 described by Hyttel, J., *Psychopharmacology* 1978, 60, 13.

Compound No.	Inhibition of Serotonin reuptake $\text{IC}_{50}$ (nM)	Compound No.	Inhibition of Serotonin reuptake $\text{IC}_{50}$ (nM)
1a	31	3b	33
1b	290	3c	46
1c	57	4a	360
1d	31	4f	14
1e	4.4	4g	23
1f	8.2	4h	71
1g	12	4l	950
1h	6.8	4m	3.0
1i	nt	4n	65
1j	360	4o	19
1k	150	4p	22
1l	nt	4q	34
2a	21	4r	12
2b	6.9	4s	24
2c	2.3	4t	99
2g	22	4u	93
2h	2.5	4v	4.4
2i	5.9	4x	nt
2t	39	5a	19
2u	20		
3a	17	Paroxetine*	0.29

Table 1 \* reference compound, nt = not tested

The 5-HT<sub>1A</sub> antagonistic activity of some of the compounds of the invention has been estimated in vitro at cloned 5-HT<sub>1A</sub> receptors stably expressed in transfected HeLa cells (HA7). In this test 5-HT<sub>1A</sub> antagonistic activity are estimated by measuring the ability of the compounds to antagonize the 5-HT induced inhibition of forskolin induced cAMP accumulation. The assay was performed as a modification of the method described by Pauwels, P.J. *et al*, *Biochem. Pharmacol.* **1993**, 45, 375.

Some of the compounds of the invention have also been tested for their *in vivo* effect on 5-HT<sub>1A</sub> receptors in the assay described by Sánchez. C. *Et al.*, *Eur. J. Pharmacol.*, **1996**, 315, pp 245. In this test antagonistic effects of test compounds are determined by measuring the ability of the test compounds to inhibit 5-MeO-DMT induced 5-HT syndrome.

The compounds of the present invention possess valuable activity as serotonin re-uptake inhibitors and have antagonistic effect at 5-HT<sub>1A</sub> receptors. The compounds of the invention are therefore considered useful for the treatment of diseases and disorders responsive to the inhibition of serotonin re-uptake and antagonistic activity at 5-HT<sub>1A</sub> receptors. Diseases responsive to the inhibition of serotonin re-uptake are well known in the art and include affective disorders, such as depression, psychosis, anxiety disorders including general anxiety disorder and panic disorder, obsessive compulsive disorder, etc.

As explained above, the antagonistic activity at 5-HT<sub>1A</sub> receptors of the compounds of the invention is predicted to counteract the negative feed back mechanism induced by the inhibition of serotonin reuptake. The antagonistic effect at 5-HT<sub>1A</sub> receptors is thus expected to improve the effect of the serotonin reuptake inhibiting activity of the compounds of the invention.

The compounds as claimed herein are therefore considered to be particularly useful as fast onset of action medicaments for the treatment of depression. The compounds may also be useful for the treatment of depressions which are non-responsive to currently available SSRI's.

**Pharmaceutical formulation**

The pharmaceutical formulations of the invention may be prepared by conventional methods in the art. For example: Tablets may be prepared by mixing the active  
5 ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvants or additives usually used for such purposes such as colourings, flavourings, preservatives etc. may be used provided that they are  
10 compatible with the active ingredients.

Solutions for injections may be prepared by dissolving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to desired volume, sterilization of the solution and filling in suitable ampules or vials. Any suitable additive conventionally used in the art may be  
15 added, such as tonicity agents, preservatives, antioxidants, etc.


The pharmaceutical compositions of this invention or those which are manufactured in accordance with this invention may be administered by any suitable route, for example orally in the form of tablets, capsules, powders, syrups, etc., or parenterally  
20 in the form of solutions for injection. For preparing such compositions, methods well known in the art may be used, and any pharmaceutically acceptable carriers, diluents, excipients, or other additives normally used in the art may be used.

Conveniently, the compounds of the invention are administered in unit dosage form containing said compounds in an amount of about 0.01 to 100 mg. The total daily  
25 dose is usually in the range of about 0.05 - 500 mg, and most preferably about 0.1 to 50 mg of the active compound of the invention.




**Claims:**

1. A substituted 4-, 5-, 6-, or 7-indole or indoline derivative of Formula



wherein W is N, C, CH or COH and the dotted lines indicate optional bonds and

wherein A is a group having the formula



wherein X is CR<sup>1A</sup>, CHR<sup>1A</sup>, N, NR<sup>1B</sup>, O, or S, where R<sup>1A</sup> is as defined for R<sup>3</sup> to R<sup>9</sup> below, and where R<sup>1B</sup> is as defined for R<sup>10</sup> below;


Y is CR<sup>2A</sup>, CHR<sup>2A</sup>, N, NR<sup>2B</sup>, O, or S, where R<sup>2A</sup> is as defined for R<sup>3</sup> to R<sup>9</sup> below and where R<sup>2B</sup> is as defined for R<sup>10</sup> below, and

the dotted lines indicate optional bonds;

provided that X and Y are not both O or S;

48

A is a group having the formula




- 5 wherein X is  $\text{CR}^{1A}$ ,  $\text{CHR}^{1A}$ , N,  $\text{NR}^{1B}$ , O, or S, where  $\text{R}^{1A}$  is as defined for  $\text{R}^3$  to  $\text{R}^9$  below, and where  $\text{R}^{1B}$  is as defined for  $\text{R}^{10}$  below;

U is C, CH, or N; and

- 10 the dotted lines indicate optional bonds; or

A is a group having the formula



15

wherein U is C, CH, or N;

Y is  $\text{CR}^{2A}$ ,  $\text{CHR}^{2A}$ , N,  $\text{NR}^{2B}$ , O, or S, where  $\text{R}^{2A}$  is as defined for  $\text{R}^3$  to  $\text{R}^9$  below and where  $\text{R}^{2B}$  is as defined for  $\text{R}^{10}$  below; and

20

the dotted lines indicate optional bonds;

n is 0, 1, 2, 3, 4, or 5, and m is 0, 1, 2, 3, 4, or 5;

Z is CH<sub>2</sub>, O, S, CO, SO, or SO<sub>2</sub>, provided that if n is 0 then Z is CH<sub>2</sub>;

- R<sup>3</sup>-R<sup>9</sup> and R<sup>11</sup> to R<sup>12</sup> are independently selected from hydrogen, halogen, cyano, nitro, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub>-alkylthio, hydroxy, hydroxy-C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxycarbonyl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, C<sub>1-6</sub>-alkylcarbonyl, phenylcarbonyl, halogen substituted phenylcarbonyl, trifluoromethyl, trifluoromethylsulfonyloxy, C<sub>1-6</sub> alkylsulfonyl, aryl and heteroaryl, and/or two adjacent groups taken from R<sup>3</sup> - R<sup>9</sup> may together form a methylenedioxy group,
- and/or two adjacent groups R<sup>7</sup> - R<sup>9</sup> may together form a cyclopentyl or cyclohexyl ring which may be substituted with one or more methyl groups, and/or one of R<sup>3</sup>-R<sup>9</sup> may alternatively be a group -NR<sup>13</sup>R<sup>14</sup> wherein R<sup>13</sup> is as defined for R<sup>10</sup> below and R<sup>14</sup> is hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub> alk(en/yn)yl, aryl, heteroaryl, aryl-C<sub>1-6</sub> alkyl, or heteroaryl-C<sub>1-6</sub>-alkyl;


R<sup>10</sup> is

- hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, aryl, heteroaryl, aryl-C<sub>1-6</sub>-alkyl, heteroaryl-C<sub>1-6</sub>-alkyl, acyl, thioacyl, C<sub>1-6</sub>-alkylsulfonyl, trifluoromethylsulfonyl, arylsulfonyl, or heteroarylsulfonyl;
- R<sup>15</sup>VCO- wherein V is O or S and R<sup>15</sup> is C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, aryl, or heteroaryl; or
- a group R<sup>16</sup>R<sup>17</sup>NCO- or R<sup>16</sup>R<sup>17</sup>NCS- wherein R<sup>16</sup> and R<sup>17</sup> are independently selected from hydrogen, C<sub>1-6</sub>-alk(en/yn)yl, C<sub>3-8</sub>-cycloalk(en)yl, C<sub>3-8</sub>-cycloalk(en)yl-C<sub>1-6</sub>-alk(en/yn)yl, heteroaryl, or aryl, or R<sup>16</sup> and R<sup>17</sup> together with the N-atom to which they are linked, form a pyrrolidinyl, piperidinyl, morpholinyl, or perhydroazepin group;

or an acid addition salt thereof.

2. A compound according to claim 1 wherein A is a group having the formula


50



wherein X, Y, the dotted lines and  $R^3$  -  $R^6$  are as defined in claim 1.

5

3. A compound according to claim 2 wherein A is a group having the formula




10

wherein  $R^3$  to  $R^6$  and the dotted lines are as defined in claim 2.

51

4. A compound according to claim 3 wherein A is a group having the formula




5

wherein  $R^3$  to  $R^6$  and the dotted lines are as defined in claim 3.

5. A compound according to claim 1 having the formula

10



(II)


wherein  $R^7$  to  $R^{12}$ ,  $W$ ,  $A$ ,  $Z$ ,  $n$ ,  $m$  and the dotted lines are as defined in claim 1.

15


6. A compound according to claims 1 to 5 wherein  $Z$  is  $CH_2$  and  $n + m$  is 0, 1, 2, 3, 4, 5, or 6.

7. A compound according to claim 1-6 wherein W is N.

8. A compound according to claim 1 having the formula









wherein  $R^7$  to  $R^{12}$ , W, Z, n, m and the dotted lines are as defined in claim 1 and A is a group having the formula




wherein X, Y, the dotted lines and  $R^3$ - $R^6$  is as defined in claim 1.

9. A compound according to claim 8 wherein A is a group having the formula

53




or



wherein  $R^3$  to  $R^6$  and the dotted line is as defined in claim 8.

5

10. A compound according to claim 9 wherein A is a group having the formula



wherein  $R^3$  to  $R^6$  and the dotted line is as defined in claim 9.

5

11. A compound of claims 1-10 wherein Z is  $\text{CH}_2$  and  $n + m$  is 0, 1, 2, 3, 4, 5, or 6 and  $R^3$ - $R^9$  and  $R^{11}$ - $R^{12}$  is hydrogen, halogen, cyano, nitro,  $\text{C}_{1-6}$ -alkyl,  $\text{C}_{1-6}$ -alkoxy hydroxy, hydroxy- $\text{C}_{1-6}$ -alkyl,  $\text{C}_{1-6}$ -alkoxycarbonyl and trifluoromethyl; and  $R^{10}$  is hydrogen.

10

12. A compound according to claim 8 -11 wherein W is N.

13. A compound according to claim 1 which is

1-(2-(3-Benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,

15 1-(3-Benzofuranylmethyl)-4-(1H-indol-4-yl)piperazine,

1-(2-(5-Fluoro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,

1-(4-(5-Fluoro-3-benzofuranyl)-1-butyl)-4-(1H-indol-4-yl)piperazine,

1-(2-(1H-Indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,

1-(3-(1H-Indol-3-yl)-1-propyl)-4-(1H-indol-4-yl)piperazine,

20 1-(4-(1H-Indol-3-yl)-1-butyl)-4-(1H-indol-4-yl)piperazine,

1-(3-(5-Fluoro-3-benzofuranyl)-1-propyl)-4-(1H-indol-4-yl)piperazine,



- 1-(2-(2-Methyl-4,5,6,7-tetrafluoro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(3-Indazolyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-3-indazolyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(7-Cyano-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
5 1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(4-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)-1,2,3,6-tetrahydropyridine,  
1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)-1,2,3,6-tetrahydropyridine,  
10 1-(2-(7-Bromo-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(1-Allyl-1H-indol-4-yl)-4-(2-(6-chloro-1H-indol-3-yl)ethyl)piperazine,  
1-(1-Allyl-1H-indol-4-yl)-4-(2-(5-fluoro-1H-indol-3-yl)ethyl)piperazine,  
1-(1-Benzyl-1H-indol-4-yl)-4-(2-(6-chloro-1H-indol-3-yl)ethyl)piperazine,  
1-(1-Benzyl-1H-indol-4-yl)-4-(2-(5-fluoro-1H-indol-3-yl)ethyl)piperazine,  
15 1-(1-Benzyl-1H-indol-4-yl)-4-(2-(5-bromo-1H-indol-3-yl)ethyl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine,  
1-(2-(1H-Indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine,  
1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine,  
1-(2-(5-Bromo-1H-indol-3-yl)ethyl)-4-(1-propargyl-1H-indol-4-yl)piperazine,  
20 1-(1-Benzyl-1H-indol-4-yl)-4-(2-(1H-indol-3-yl)ethyl)piperazine,  
1-(2-(5-Bromo-1H-indol-3-yl)ethyl)-4-(1H-indol-5-yl)piperazine,  
1-(2-(5-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-5-yl)piperazine,  
1-(2-(5-Fluoro-1H-indol-3-yl)ethyl)-4-(6-hydroxymethyl-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-hydroxymethyl-1H-indol-4-yl)piperazine,  
25 1-(2-(5-Bromo-1H-indol-3-yl)ethyl)-4-(6-hydroxymethyl-1H-indol-4-yl)piperazine,  
1-(3-(6-Fluoro-1,2-benzisoxazol-3-yl)-1-propyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(1H-Indol-3-yl)ethyl)-4-(6-methoxycarbonyl-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-methoxycarbonyl-1H-indol-4-yl)piperazine,  
1-(2-(5-Fluoro-3-benzofuranyl)ethyl)-4-(6-methoxycarbonyl-1H-indol-4-  
30 yl)piperazine,  
1-(5-Fluoro-3-benzofuranylmethyl)-4-(1H-indol-4-yl)piperazine,  
1-(3-Cyano-1H-indol-4-yl)-4-(2-(1H-indol-3-yl)ethyl)piperazine,

- 1-(3-Cyano-1H-indol-4-yl)-4-(2-(5-fluoro-3-benzofuranyl)ethyl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(3-cyano-1H-indol-4-yl)piperazine,  
1-(2-(3-Benzofuranyl)ethyl)-4-(3-cyano-1H-indol-4-yl)piperazine,  
1-(1H-Indol-4-yl)-4-(2-(5-methyl-3-benzofuranyl)ethyl)piperazine,  
5 1-(1H-Indol-4-yl)-4-(2-(4-methyl-3-benzofuranyl)ethyl)piperazine,  
1-(3-(5-Fluoro-3-benzofuranyl)-1-propyl)-4-(1H-indol-4-yl)-1,2,3,6-  
tetrahydropyridine,  
1-(2-(5-Chloro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(1H-Indol-4-yl)-4-(2-(6-methyl-3-benzofuranyl)ethyl)piperazine,  
10 1-(2-(7-Chloro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(4-Chloro-1H-indol-3-yl)ethyl)-4-(3-cyano-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)-4-(1H-indol-4-yl)piperidine,  
1-(2-(5-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(7-Bromo-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
15 1-(2-(4-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(6-Trifluoromethyl-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(1H-Indol-4-yl)-4-(2-(5-methyl-1H-indol-3-yl)ethyl)piperazine,  
1-(1H-Indol-4-yl)-4-(2-(6-methyl-1H-indol-3-yl)ethyl)piperazine,  
1-(1H-Indol-4-yl)-4-(2-(7-methyl-1H-indol-3-yl)ethyl)piperazine,  
20 1-(2-(4,5-Dichloro-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(5-Bromo-3-benzofuranyl)ethyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(4-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-4-yl)piperidine,  
4-(1H-Indol-4-yl)-1-(2-(5-methyl-1H-indol-3-yl)ethyl)piperidine,  
4-(1H-Indol-4-yl)-1-(2-(1H-indol-3-yl)ethyl)piperidine,  
25 1-(1H-Indol-4-yl)-4-(3-(4-methyl-3-benzofuranyl)-1-propyl)piperazine,  
4-(1H-Indol-4-yl)-1-(3-(4-methyl-3-benzofuranyl)-1-propyl)piperidine,  
1-(3-(4-Chloro-3-benzofuranyl)-1-propyl)-4-(1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-chloro-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-fluoro-1H-indol-4-yl)piperazine,  
30 1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(6-cyano-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(7-chloro-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(7-cyano-1H-indol-4-yl)piperazine,

1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(2-cyano-1H-indol-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indolin-4-yl)piperazine,  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-6-yl)piperazine and  
1-(2-(6-Chloro-1H-indol-3-yl)ethyl)-4-(1H-indol-7-yl)piperazine or a  
5 pharmaceutically acceptable acid addition salt thereof.

14. A pharmaceutical composition comprising a compound according to claims 1  
to 13 or a pharmaceutically acceptable acid addition salt thereof and at least one  
pharmaceutically acceptable carrier or diluent.

10

15. The use of a compound according to claims 1 to 13 or a pharmaceutically  
acceptable acid addition salt thereof for the preparation of a medicament for the  
treatment of a disorder or disease responsive to the inhibition of serotonin reuptake  
and antagonism of 5-HT<sub>1A</sub> receptors.

15

16. The use of a compound according to claims 1 to 13 or a pharmaceutically  
acceptable acid addition salt thereof for the preparation of a medicament for the  
treatment of affective disorders, such as depression, psychosis, anxiety disorders  
including general anxiety disorder and panic disorder and obsessive compulsive  
20 disorder.

17. A method for the treatment of a disorder or disease of living animal body,  
including a human, which is responsive to the inhibition of serotonin reuptake and  
antagonism of 5-HT<sub>1A</sub> receptors comprising administering to such a living animal  
25 body, including a human, a therapeutically effective amount of a compound according  
to claims 1 to 13 or a pharmaceutically acceptable acid addition salt thereof.

18. A method for the treatment of an affective disorder, including depression  
psychosis, anxiety disorders including general anxiety disorder and panic disorder and  
30 obsessive compulsive disorder in a living animal body, including a human,  
comprising administering a therapeutically effective amount of a compound according  
to claims 1 to 13 or a pharmaceutically acceptable acid addition salt thereof.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 99/00326

## A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07D 405/14, C07D 405/12, C07D 401/14, C07D 209/20, A61K 31/495,  
A61K 31/445

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9421626 A1 (MERCK SHARP & DOHME LIMITED), 29 Sept 1994 (29.09.94), examples 19 and 20 --	1-16
P,X	WO 9828293 A1 (H. LUNDBECK A/S), 2 July 1998 (02.07.98) --	1,5-7,11-12, 14-16
A	European Journal of Pharmacology - Molecular Pharmacology Section, Volume 244, 1993, Philippe Schoeffter et al, "SDZ 216-525, a selective and potent 5-HT1A receptor antagonist" page 251 - page 257 --	1-16

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

6 Sept 1999

Date of mailing of the international search report

16-10-1999

Name and mailing address of the ISA/  
Swedish Patent Office  
Box 5055, S-102 42 STOCKHOLM  
Facsimile No. +46 8 666 02 86

Authorized officer

Göran Karlsson/EÖ  
Telephone No. +46 8 782 25 00

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 99/00326

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Pharmacology Biochemistry and Behavior, Volume 46, 1993, Robert Bell et al, "Effects of (-)-Pindolol and SDZ 216-525 on Social and Agonistic Behavior in Mice" page 873 - page 880 --	1-16
A	DE 4414113 A1 (MERCK PATENT GMBH), 26 October 1995 (26.10.95) -- -----	1-16

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/DK99/00326

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 17-18  
because they relate to subject matter not required to be searched by this Authority, namely:  
**A method for treatment of the human or animal body by therapy,  
see rule 39.1**
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/DK 99/00326

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9421626	A1	29/09/94	AU	679214 B	26/06/97
				AU	6213994 A	11/10/94
				CA	2156839 A	29/09/94
				EP	0701555 A	20/03/96
				JP	8507780 T	20/08/96
				US	5665732 A	09/09/97
-----						
WO	9828293	A1	02/07/98	AU	5310998 A	17/07/98
				ZA	9711376 A	21/07/98
-----						
DE	4414113	A1	26/10/95	AT	172730 T	15/11/98
				AU	697749 B	15/10/98
				AU	1648895 A	02/11/95
				CA	2147451 A	23/10/95
				CN	1114651 A	10/01/96
				CZ	9501035 A	13/12/95
				DE	59504032 D	00/00/00
				EP	0683166 A,B	22/11/95
				SE	0683166 T3	
				ES	2125508 T	01/03/99
				HU	74096 A	28/11/96
				HU	9501139 D	00/00/00
				JP	7291969 A	07/11/95
				NO	951529 A	23/10/95
				PL	308287 A	30/10/95
				SK	50895 A	08/11/95
				US	5693655 A	02/12/97
				ZA	9503260 A	09/01/96
-----						

**THIS PAGE BLANK (USPTO)**